

GKZ hypergeometric systems of the four-loop vacuum Feynman integrals

Hai-Bin Zhang^{1,2,3,4,*}, Tai-Fu Feng^{1,2,3,5,†}

¹*Department of Physics, Hebei University, Baoding, 071002, China*

²*Hebei Key Laboratory of High-precision Computation and Application of Quantum Field Theory, Baoding, 071002, China*

³*Hebei Research Center of the Basic Discipline for Computational Physics, Baoding, 071002, China*

⁴*Institute of Life Science and Green Development, Hebei University, Baoding, 071002, China*

⁵*Department of Physics, Chongqing University, Chongqing, 401331, China*

Abstract

Basing on Mellin-Barnes representations and Miller's transformation, we present the Gel'fand-Kapranov-Zelevinsky (GKZ) hypergeometric systems of the four-loop vacuum Feynman integrals with arbitrary masses. Through the GKZ hypergeometric systems, the analytical hypergeometric series solutions of the four-loop vacuum Feynman integrals with arbitrary masses can be obtained in neighborhoods of origin including infinity. The analytical expressions of the four-loop vacuum Feynman integrals can be formulated as a linear combination of the corresponding fundamental solution systems in certain convergent region.

PACS numbers: 02.30.Jr, 11.10.Gh, 12.38.Bx

Keywords: Four-loop vacuum Feynman integral, Linear partial differential equation, GKZ hypergeometric system, analytical calculation

* hbzhang@hbu.edu.cn

† fengtf@hbu.edu.cn

Contents

I. Introduction	3
II. GKZ hypergeometric system of the four-loop vacuum integral with five propagates	4
III. GKZ hypergeometric system of the four-loop vacuum integrals with six propagates	9
A. Four-loop vacuum diagram with six propagates for type A	9
B. Four-loop vacuum diagram with six propagates for type B	12
IV. GKZ hypergeometric system of the four-loop vacuum integrals with seven propagates	15
A. Four-loop vacuum diagram with seven propagates for type A	15
B. Four-loop vacuum diagram with seven propagates for type B	19
V. The hypergeometric series solutions of the four-loop vacuum integrals	22
A. The general case of the four-loop vacuum integral with five propagates	22
B. The special case of the four-loop vacuum integral with five propagates	26
C. The four-loop vacuum integrals with six or seven propagates	29
VI. Conclusions	30
Acknowledgments	31
A. The calculation of the integral I_q	31
B. The hypergeometric series solutions of the integer lattice B_{1345}	32
C. The hypergeometric series solutions of the integer lattice $B_{\bar{1}345}$	36
References	40

I. INTRODUCTION

With the improvement of experimental measurement accuracy at the planned future colliders [1–6], Feynman integrals need to be calculated beyond two-loop order. Vacuum integrals are the important subsets of Feynman integrals, which constitute a main building block in asymptotic expansions of Feynman integrals [7, 8] and are also useful for massless theories when a propagator mass is introduced as an intermediate infrared regulator [9]. The calculation of multi-loop vacuum integrals is a good breakthrough window in the calculation of multi-loop Feynman integrals. In this article, we investigate the analytical calculation for the four-loop vacuum integrals with arbitrary masses.

It's well known that the completely general one-loop integrals are analytically in the time-space dimension $D = 4 - 2\epsilon$ [10–13]. At the two-loop level, the vacuum integrals have been calculated to polylogarithms or equivalent functions [14–18]. The three-loop vacuum integrals are also calculated analytically and numerically in some literatures [19–40]. But, very few four-loop vacuum integrals are calculated analytically. The vacuum integrals at the four-loop level are only calculated analytically for single-mass-scale [41, 42], equal masses [43], and reduction [44, 45]. Recently, Feynman integrals using auxiliary mass flow numerical method [46, 47], also can be reduced to vacuum integrals, which can be numerical solved by further reduction. In order to further improve the computational efficiency and give analytical results completely, it is meaningful to explore new analytical calculating method of the four-loop vacuum integrals with arbitrary masses.

During the past decades, Feynman integrals have been considered as the generalized hypergeometric functions [48–72]. Considering Feynman integrals as the generalized hypergeometric functions, one finds that the D -module of a Feynman diagram [64, 73] is isomorphic to Gel'fand-Kapranov-Zelevinsky (GKZ) D -module [74–78]. GKZ-hypergeometric systems of Feynman integrals with codimension= 0, 1 are presented in Refs. [79, 80] through Lee-Pomeransky parametric representations [81]. To construct canonical series solutions with suitable independent variables, one should compute the restricted D -module of GKZ-hypergeometric system originating from Lee-Pomeransky representations on corresponding hyperplane in the parameter space [82–84]. In our previous work, GKZ hypergeometric systems of one- and two-loop Feynman diagrams are also obtained [85–87] from Mellin-Barnes representations [68, 69], through Miller's transformation [88, 89]. There are some recent

work in GKZ framework of Feynman integrals [90–108].

In our previous work, we have given GKZ hypergeometric systems of the Feynman integrals of the two-loop vacuum integral [85] and three-loop vacuum integrals [40]. In this article, we derive GKZ hypergeometric systems of the four-loop vacuum integrals with arbitrary masses, basing on Mellin-Barnes representations and Miller’s transformation. The generally strategy for analyzing the four-loop vacuum integrals includes three steps here. Firstly, we obtain the Mellin-Barnes representation of the vacuum integral. Secondly, we find GKZ hypergeometric system of the vacuum integral via Miller’s transformation. Finally, analytical hypergeometric series solutions of the vacuum integral are constructed in neighborhoods of origin including infinity. The integration constants, i.e. the combination coefficients, are determined from the vacuum integral of an ordinary point or some regular singularities.

Our presentation is organized as following. Through the Mellin-Barnes representation and Miller’s transformation, we derive the GKZ hypergeometric system of the four-loop vacuum integral with five propagates in Sec. II, six propagates in Sec. III, and seven propagates in Sec. IV. And then, we construct the analytical hypergeometric series solutions of the GKZ system of the four-loop vacuum integrals in Sec. V. At last, the conclusions are summarized in Sec. VI, and some formulates are presented in the appendices.

II. GKZ HYPERGEOMETRIC SYSTEM OF THE FOUR-LOOP VACUUM INTEGRAL WITH FIVE PROPAGATES

The general analytic expression for the Feynman integral of the four-loop vacuum diagram with five propagates in Fig. 1 is written as

$$U_5 = \left(\Lambda_{\text{RE}}^2\right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2)(q_2^2 - m_2^2)(q_3^2 - m_3^2)} \times \frac{1}{[(q_1 + q_2 + q_3 + q_4)^2 - m_4^2](q_4^2 - m_5^2)}, \quad (1)$$

where $D = 4 - 2\varepsilon$ is the number of dimensions in dimensional regularization and Λ_{RE} denotes the renormalization energy scale, $\mathbf{q} = (q_1, \dots, q_4)$, $\frac{d^D \mathbf{q}}{(2\pi)^D} = \frac{d^D q_1}{(2\pi)^D} \frac{d^D q_2}{(2\pi)^D} \frac{d^D q_3}{(2\pi)^D} \frac{d^D q_4}{(2\pi)^D}$. The Feynman integral is hard to calculate analytically, if all virtual masses are nonzero. So, one can extract the virtual masses from the integral, to facilitate further calculation. Adopting




FIG. 1: Four-loop vacuum diagram with five propagators, which m_i denotes the mass of the i -th particle and q_j denotes the momentum.

the notation of Refs. [68, 69], the Feynman integral of the four-loop vacuum diagram with five propagates can be written as

$$U_5 = \frac{(\Lambda_{\text{RE}}^2)^{8-2D}}{(2\pi i)^4} \int_{-i\infty}^{+i\infty} d\mathbf{s} \left[\prod_{i=1}^4 (-m_i^2)^{s_i} \Gamma(-s_i) \Gamma(1+s_i) \right] I_q, \quad (2)$$

where $\mathbf{s} = (s_1, \dots, s_4)$, $d\mathbf{s} = ds_1 ds_2 ds_3 ds_4$ and

$$I_q \equiv \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2)^{1+s_1} (q_2^2)^{1+s_2} (q_3^2)^{1+s_3} [(q_1 + q_2 + q_3 + q_4)^2]^{1+s_4} (q_4^2 - m_5^2)}. \quad (3)$$

The integral I_q just keeps one mass m_5 , which can be easily calculated analytically. The calculation of the integral I_q can be seen in Appendix A.

And then, the Mellin-Barnes representation of the Feynman integral of the four-loop vacuum diagram in Eq. (2) can be written as

$$U_5 = \frac{-m_5^6}{(2\pi i)^4 (4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_5^2} \right)^{8-2D} \int_{-i\infty}^{+i\infty} d\mathbf{s} \left[\prod_{i=1}^4 \left(\frac{m_i^2}{m_5^2} \right)^{s_i} \Gamma(-s_i) \right] \\ \times \left[\prod_{i=1}^4 \Gamma\left(\frac{D}{2} - 1 - s_i\right) \right] \Gamma\left(4 - \frac{3D}{2} + \sum_{i=1}^4 s_i\right) \Gamma\left(5 - 2D + \sum_{i=1}^4 s_i\right). \quad (4)$$

It is well known that negative integers and zero are simple poles of the function $\Gamma(z)$. As all s_i contours are closed to the right in corresponding complex planes, one finds that the analytic expression of the Feynman integral can be written as the linear combination of generalized hypergeometric functions. Taking the residue of the pole of $\Gamma(-s_i)$, ($i =$

$1, \dots, 4)$, we can derive one linear independent term of the integral:

$$U_5 \supset \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2} \right)^{8-2D} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \sum_{n_4=0}^{\infty} (-)^{\sum_{i=1}^4 n_i} x_1^{n_1} x_2^{n_2} x_3^{n_3} x_4^{n_4} \\ \times \left[\prod_{i=1}^4 \Gamma\left(\frac{D}{2} - 1 - n_i\right) (n_i!)^{-1} \right] \Gamma\left(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i\right) \Gamma\left(5 - 2D + \sum_{i=1}^4 n_i\right), \quad (5)$$

with $x_i = \frac{m_i^2}{m_5^2}$, ($i = 1, \dots, 4$).

We adopt the identity

$$\Gamma(z - n) \Gamma(1 - z + n) = (-)^n \Gamma(z) \Gamma(1 - z) = (-)^n \pi / \sin \pi z, \quad (6)$$

which originate from the well-known relation $\Gamma(z) \Gamma(1 - z) = \pi / \sin \pi z$. Then, Eq. (5) can be written as

$$U_5 \supset \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2} \right)^{8-2D} \frac{\pi^4}{\sin^4 \frac{\pi D}{2}} T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (7)$$

with

$$T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \sum_{n_4=0}^{\infty} A_{n_1 n_2 n_3 n_4} x_1^{n_1} x_2^{n_2} x_3^{n_3} x_4^{n_4}, \quad (8)$$

where $\mathbf{x} = (x_1, x_2, x_3, x_4)$, $\mathbf{a} = (a_1, a_2)$ and $\mathbf{b} = (b_1, b_2, b_3, b_4)$ with

$$a_1 = 4 - \frac{3D}{2}, \quad a_2 = 5 - 2D, \quad b_1 = b_2 = b_3 = b_4 = 2 - \frac{D}{2}, \quad (9)$$

and the coefficient $A_{n_1 n_2 n_3 n_4}$ is

$$A_{n_1 n_2 n_3 n_4} = \frac{\Gamma(a_1 + \sum_{i=1}^4 n_i) \Gamma(a_2 + \sum_{i=1}^4 n_i)}{\prod_{i=1}^4 n_i! \Gamma(b_i + n_i)}. \quad (10)$$

Here, we just derive one linear independent term of the integral. We still need to derive other linear independent terms of the integral.

Through the adjacent relations of the coefficient $A_{n_1 n_2 n_3 n_4}$:

$$\frac{A_{(n_1+1)n_2 n_3 n_4}}{A_{n_1 n_2 n_3 n_4}} = \frac{(a_1 + \sum_{i=1}^4 n_i)(a_2 + \sum_{i=1}^4 n_i)}{(n_1 + 1)(b_1 + n_1)}, \\ \frac{A_{n_1(n_2+1)n_3 n_4}}{A_{n_1 n_2 n_3 n_4}} = \frac{(a_1 + \sum_{i=1}^4 n_i)(a_2 + \sum_{i=1}^4 n_i)}{(n_2 + 1)(b_2 + n_2)},$$

$$\begin{aligned}\frac{A_{n_1 n_2 (n_3+1) n_4}}{A_{n_1 n_2 n_3 n_4}} &= \frac{(a_1 + \sum_{i=1}^4 n_i)(a_2 + \sum_{i=1}^4 n_i)}{(n_3 + 1)(b_3 + n_3)}, \\ \frac{A_{n_1 n_2 n_3 (n_4+1)}}{A_{n_1 n_2 n_3 n_4}} &= \frac{(a_1 + \sum_{i=1}^4 n_i)(a_2 + \sum_{i=1}^4 n_i)}{(n_4 + 1)(b_4 + n_4)},\end{aligned}\tag{11}$$

the difference-differential operators are written as

$$\begin{aligned}(\sum_{i=1}^4 \vartheta_{x_i} + a_j)T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) &= a_j T_5(\mathbf{a} + \mathbf{e}_{2,j}, \mathbf{b} \mid \mathbf{x}), \quad (j = 1, 2), \\ (\vartheta_{x_k} + b_k - 1)T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) &= (b_k - 1)T_5(\mathbf{a}, \mathbf{b} - \mathbf{e}_{4,k} \mid \mathbf{x}), \\ \partial_{x_k} T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) &= \frac{a_1 a_2}{b_k} T_5(\mathbf{a} + \mathbf{e}_2, \mathbf{b} + \mathbf{e}_{4,k} \mid \mathbf{x}), \quad (k = 1, \dots, 4).\end{aligned}\tag{12}$$

Here $\mathbf{e}_{2,j} \in \mathbf{R}^2$ ($j = 1, 2$) with $\mathbf{e}_{2,1} = (1, 0)$ and $\mathbf{e}_{2,2} = (0, 1)$, $\mathbf{e}_2 = (1, 1)$, $\mathbf{e}_{4,k} \in \mathbf{R}^4$ ($k = 1, \dots, 4$) denotes the row vector whose entry is zero except that the k -th entry is 1, $\mathbf{e}_4 = (1, 1, 1, 1)$, $\vartheta_{x_k} = x_k \partial_{x_k}$ denotes the Euler operators, and $\partial_{x_k} = \partial/\partial x_k$, respectively.

We can define the auxiliary function

$$\Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = \mathbf{u}^{\mathbf{a}} \mathbf{v}^{\mathbf{b} - \mathbf{e}_4} T_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}),\tag{13}$$

with the intermediate variables $\mathbf{u} = (u_1, u_2) = (1, 1)$, $\mathbf{v} = (v_1, v_2, v_3, v_4) = (1, 1, 1, 1)$. Through Miller's transformation [88, 89], the relations are obtained

$$\begin{aligned}\vartheta_{u_j} \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= a_j \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (j = 1, 2), \\ \vartheta_{v_k} \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= (b_k - 1) \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (k = 1, \dots, 4),\end{aligned}\tag{14}$$

which naturally induces the notion of GKZ hypergeometric system.

In addition, the contiguous relations of Eq. (12) are rewritten as

$$\begin{aligned}u_j (\sum_{i=1}^4 \vartheta_{x_i} + \vartheta_{u_j}) \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= a_j \Phi_5(\mathbf{a} + \mathbf{e}_{2,j}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \\ \frac{1}{v_k} (\vartheta_{x_k} + \vartheta_{v_k}) \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= (b_k - 1) \Phi_5(\mathbf{a}, \mathbf{b} - \mathbf{e}_{4,k} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \\ u_1 u_2 v_k \partial_{x_k} \Phi_5(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= \frac{a_1 a_2}{b_k} \Phi_5(\mathbf{a} + \mathbf{e}_2, \mathbf{b} + \mathbf{e}_{4,k} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}).\end{aligned}\tag{15}$$

The operators above together with ϑ_{u_j} , ϑ_{v_j} define the Lie algebra of the hypergeometric systems [88, 89]. Through the transformation

$$z_j = \frac{1}{u_j}, \quad (j = 1, 2), \quad z_{2+k} = v_k, \quad z_{6+k} = \frac{x_k}{u_1 u_2 v_k}, \quad (k = 1, \dots, 4),\tag{16}$$

one derives

$$\begin{aligned}
\vartheta_{u_1} &= -\vartheta_{z_1} - \vartheta_{z_7} - \vartheta_{z_8} - \vartheta_{z_9} - \vartheta_{z_{10}} , \\
\vartheta_{u_2} &= -\vartheta_{z_2} - \vartheta_{z_7} - \vartheta_{z_8} - \vartheta_{z_9} - \vartheta_{z_{10}} , \\
\vartheta_{v_k} &= \vartheta_{z_{2+k}} - \vartheta_{z_{6+k}} , \quad \vartheta_{x_k} = \vartheta_{z_{6+k}} .
\end{aligned} \tag{17}$$

Through Eq. (14), one can finally have the GKZ hypergeometric system for the four-loop vacuum integral with five propagates:

$$\mathbf{A}_5 \cdot \vec{\vartheta}_5 \Phi_5 = \mathbf{B}_5 \Phi_5 , \tag{18}$$

where

$$\begin{aligned}
\mathbf{A}_5 &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 \end{pmatrix} , \\
\vec{\vartheta}_5^T &= (\vartheta_{z_1}, \dots, \vartheta_{z_{10}}) , \\
\mathbf{B}_5^T &= (-a_1, -a_2, b_1 - 1, b_2 - 1, b_3 - 1, b_4 - 1) .
\end{aligned} \tag{19}$$

Correspondingly the dual matrix $\tilde{\mathbf{A}}_5$ of \mathbf{A}_5 is

$$\tilde{\mathbf{A}}_5 = \begin{pmatrix} -1 & -1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{pmatrix} . \tag{20}$$

The row vectors of the matrix $\tilde{\mathbf{A}}_5$ induce the integer sublattice \mathbf{B} which can be used to construct the formal solutions in hypergeometric series.

Defining the combined variables

$$y_1 = \frac{z_3 z_7}{z_1 z_2}, \quad y_2 = \frac{z_4 z_8}{z_1 z_2}, \quad y_3 = \frac{z_5 z_9}{z_1 z_2}, \quad y_4 = \frac{z_6 z_{10}}{z_1 z_2}, \tag{21}$$

we write the solutions satisfying Eq. (18) as

$$\Phi_5(\mathbf{z}) = \left(\prod_{i=1}^{10} z_i^{\alpha_i} \right) \varphi_5(y_1, y_2, y_3, y_4) . \tag{22}$$




FIG. 2: Four-loop vacuum diagram with six propagators for type A, which m_i denotes the mass of the i -th particle and q_j denotes the momentum.

Here $\vec{\alpha}^T = (\alpha_1, \dots, \alpha_{10})$ denotes a sequence of complex number such that

$$\mathbf{A}_5 \cdot \vec{\alpha} = \mathbf{B}_5, \quad (23)$$

namely,

$$\begin{aligned} \alpha_1 + \alpha_7 + \alpha_8 + \alpha_9 + \alpha_{10} &= -a_1, & \alpha_2 + \alpha_7 + \alpha_8 + \alpha_9 + \alpha_{10} &= -a_2, \\ \alpha_3 - \alpha_7 &= b_1 - 1, & \alpha_4 - \alpha_8 &= b_2 - 1, & \alpha_5 - \alpha_9 &= b_3 - 1, & \alpha_6 - \alpha_{10} &= b_4 - 1. \end{aligned} \quad (24)$$

In the following Sec. V, we will show the analytical hypergeometric series solutions solving from the GKZ hypergeometric system in Eq. (18).

III. GKZ HYPERGEOMETRIC SYSTEM OF THE FOUR-LOOP VACUUM INTEGRALS WITH SIX PROPAGATES

A. Four-loop vacuum diagram with six propagates for type A

The four-loop vacuum diagrams with six propagates have two topologies, which one can be seen in Fig. 2. The general analytic expression for the Feynman integral of the four-loop vacuum diagram with six propagates for type A in Fig. 2 is written as

$$\begin{aligned} U_{6A} &= \left(\Lambda_{\text{RE}}^2 \right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2)(q_2^2 - m_2^2)[(q_1 + q_2 + q_3 + q_4)^2 - m_3^2]} \\ &\quad \times \frac{1}{[(q_3 + q_4)^2 - m_4^2](q_3^2 - m_5^2)(q_4^2 - m_6^2)}. \end{aligned} \quad (25)$$

Through the Mellin-Barnes transformation, the Feynman integral can be written as

$$U_{6A} = \frac{(\Lambda_{\text{RE}}^2)^{8-2D}}{(2\pi i)^5} \int_{-i\infty}^{+i\infty} d\mathbf{s} \left[\prod_{i=1}^5 (-m_i^2)^{s_i} \Gamma(-s_i) \Gamma(1+s_i) \right] I_{6A}, \quad (26)$$

where $\mathbf{s} = (s_1, \dots, s_5)$, and

$$I_{6A} \equiv \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2)^{1+s_1} (q_2^2)^{1+s_2} [(q_1 + q_2 + q_3 + q_4)^2]^{1+s_3}} \times \frac{1}{[(q_3 + q_4)^2]^{1+s_4} (q_5^2)^{1+s_5} (q_6^2 - m_6^2)}. \quad (27)$$

Similarly, integrating out \mathbf{q} , the Mellin-Barnes representation of the Feynman integral of the four-loop vacuum diagram with six propagates for type A can be written as

$$U_{6A} = \frac{m_6^4}{(2\pi i)^5 (4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_6^2} \right)^{8-2D} \int_{-i\infty}^{+i\infty} d\mathbf{s} \left[\prod_{i=1}^5 \left(\frac{m_i^2}{m_6^2} \right)^{s_i} \Gamma(-s_i) \right] \times \left[\prod_{i=1}^3 \Gamma\left(\frac{D}{2} - 1 - s_i\right) \right] \Gamma\left(\frac{D}{2} - 1 - s_5\right) \Gamma\left(5 - \frac{3D}{2} + \sum_{i=1}^5 s_i\right) \times \frac{\Gamma(6 - 2D + \sum_{i=1}^5 s_i) \Gamma(3 - D + \sum_{i=1}^3 s_i) \Gamma\left(\frac{3D}{2} - 4 - \sum_{i=1}^4 s_i\right) \Gamma(1 + s_4)}{\Gamma(4 - D + \sum_{i=1}^4 s_i) \Gamma\left(\frac{3D}{2} - 3 - \sum_{i=1}^3 s_i\right)}. \quad (28)$$

Taking the residue of the pole of $\Gamma(-s_i)$, ($i = 1, \dots, 5$), one can derive one linear independent term of the vacuum integral:

$$U_{6A} \ni \frac{-m_6^4}{(4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_6^2} \right)^{8-2D} \sum_{\mathbf{n}=0}^{\infty} \mathbf{x}^{\mathbf{n}} (-)^{\sum_{i=1}^5 n_i} \left[\prod_{i=1}^3 \Gamma\left(\frac{D}{2} - 1 - n_i\right) \right] \times \Gamma\left(\frac{D}{2} - 1 - n_5\right) \Gamma\left(5 - \frac{3D}{2} + \sum_{i=1}^5 n_i\right) \Gamma(6 - 2D + \sum_{i=1}^5 n_i) \times \frac{\Gamma(3 - D + \sum_{i=1}^3 n_i) \Gamma\left(\frac{3D}{2} - 4 - \sum_{i=1}^4 n_i\right) \Gamma(1 + n_4)}{\left[\prod_{i=1}^5 n_i! \right] \Gamma(4 - D + \sum_{i=1}^4 n_i) \Gamma\left(\frac{3D}{2} - 3 - \sum_{i=1}^3 n_i\right)}, \quad (29)$$

with $\mathbf{n} = (n_1, \dots, n_5)$, $\mathbf{x} = (x_1, \dots, x_5)$, $x_i = \frac{m_i^2}{m_6^2}$, ($i = 1, \dots, 5$). Adopting the identity in Eq. (6), Eq. (29) can be written as

$$U_{6A} \ni \frac{m_6^4}{(4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_6^2} \right)^{8-2D} \frac{\pi^4}{\sin^4 \frac{\pi D}{2}} T_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (30)$$

with

$$T_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) = \sum_{\mathbf{n}=0}^{\infty} A_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}, \quad (31)$$

where $\mathbf{a} = (a_1, \dots, a_5)$ and $\mathbf{b} = (b_1, \dots, b_6)$ with

$$\begin{aligned} a_1 &= 5 - \frac{3D}{2}, \quad a_2 = 6 - 2D, \quad a_3 = 3 - D, \quad a_4 = 4 - \frac{3D}{2}, \quad a_5 = 1, \\ b_1 &= b_2 = b_3 = b_4 = 2 - \frac{D}{2}, \quad b_5 = 4 - D, \quad b_6 = 5 - \frac{3D}{2}, \end{aligned} \quad (32)$$

and the coefficient $A_{\mathbf{n}}$ is

$$A_{\mathbf{n}} = \frac{\Gamma(a_1 + \sum_{i=1}^5 n_i) \Gamma(a_2 + \sum_{i=1}^5 n_i) \Gamma(a_3 + \sum_{i=1}^3 n_i) \Gamma(a_4 + \sum_{i=1}^3 n_i) \Gamma(a_5 + n_4)}{\left[\prod_{i=1}^5 n_i! \right] \left[\prod_{i=1}^3 \Gamma(b_i + n_i) \right] \Gamma(b_4 + n_5) \Gamma(b_5 + \sum_{i=1}^4 n_i) \Gamma(b_6 + \sum_{i=1}^4 n_i)}. \quad (33)$$

In order to proceed with our analysis, we also define the auxiliary function

$$\Phi_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = \mathbf{u}^{\mathbf{a}} \mathbf{v}^{\mathbf{b} - \mathbf{e}_6} T_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (34)$$

with the intermediate variables $\mathbf{u} = (u_1, \dots, u_5) = (1, 1, 1, 1, 1)$, $\mathbf{v} = (v_1, \dots, v_6)$, $\mathbf{v} = \mathbf{e}_6 = (1, 1, 1, 1, 1, 1)$. Then one can obtain

$$\begin{aligned} \vartheta_{u_j} \Phi_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= a_j \Phi_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (j = 1, \dots, 5), \\ \vartheta_{v_k} \Phi_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= (b_k - 1) \Phi_{6A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (k = 1, \dots, 6). \end{aligned} \quad (35)$$

Through the transformation

$$\begin{aligned} z_j &= \frac{1}{u_j}, \quad (j = 1, \dots, 5), \quad z_{5+k} = v_k, \quad (k = 1, \dots, 6), \\ z_{12} &= \frac{x_1}{u_1 u_2 u_3 u_4 v_1 v_5 v_6}, \quad z_{13} = \frac{x_2}{u_1 u_2 u_3 u_4 v_2 v_5 v_6}, \\ z_{14} &= \frac{x_3}{u_1 u_2 u_3 u_4 v_3 v_5 v_6}, \quad z_{15} = \frac{x_4}{u_1 u_2 u_5 v_5 v_6}, \quad z_{16} = \frac{x_5}{u_1 u_2 v_4}, \end{aligned} \quad (36)$$

one derives the GKZ hypergeometric system for the four-loop vacuum diagram with six propagates for type A:

$$\mathbf{A}_{6A} \cdot \vec{\vartheta}_{6A} \Phi_{6A} = \mathbf{B}_{6A} \Phi_{6A}, \quad (37)$$

where

$$\mathbf{A}_{6A} = \begin{pmatrix} \mathbf{I}_{11 \times 11} & \mathbf{A}_{\mathbf{x}6A} \end{pmatrix},$$

$$\begin{aligned}
\mathbf{A}_{\mathbf{x6A}}^T &= \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & -1 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & -1 & 0 & 0 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}, \\
\vec{\vartheta}_{6A}^T &= (\vartheta_{z_1}, \dots, \vartheta_{z_{16}}), \\
\mathbf{B}_{6A}^T &= (-a_1, \dots, -a_5, b_1 - 1, \dots, b_6 - 1).
\end{aligned} \tag{38}$$

Here, $\mathbf{I}_{11 \times 11}$ is an 11×11 unit matrix.

Correspondingly the dual matrix $\tilde{\mathbf{A}}_{6A}$ of \mathbf{A}_{6A} is

$$\tilde{\mathbf{A}}_{6A} = \begin{pmatrix} -\mathbf{A}_{\mathbf{x6A}}^T & \mathbf{I}_{5 \times 5} \end{pmatrix}. \tag{39}$$

Here, $\mathbf{I}_{5 \times 5}$ is a 5×5 unit matrix. The row vectors of the matrix $\tilde{\mathbf{A}}_{6A}$ induce the integer sublattice \mathbf{B} which can be used to construct the formal solutions in hypergeometric series.

Defining the combined variables

$$\begin{aligned}
y_1 &= \frac{z_6 z_{10} z_{11} z_{12}}{z_1 z_2 z_3 z_4}, & y_2 &= \frac{z_7 z_{10} z_{11} z_{13}}{z_1 z_2 z_3 z_4}, \\
y_3 &= \frac{z_8 z_{10} z_{11} z_{14}}{z_1 z_2 z_3 z_4}, & y_4 &= \frac{z_{10} z_{11} z_{15}}{z_1 z_2 z_5}, & y_5 &= \frac{z_9 z_{16}}{z_1 z_2},
\end{aligned} \tag{40}$$

we write the solutions satisfying Eq. (37) as

$$\Phi_{6A}(\mathbf{z}) = \left(\prod_{i=1}^{16} z_i^{\alpha_i} \right) \varphi_{6A}(\mathbf{y}), \tag{41}$$

where $\mathbf{y} = (y_1, \dots, y_5)$, $\vec{\alpha}^T = (\alpha_1, \dots, \alpha_{16})$ denotes a sequence of complex number such that

$$\mathbf{A}_{6A} \cdot \vec{\alpha} = \mathbf{B}_{6A}. \tag{42}$$

B. Four-loop vacuum diagram with six propagates for type B

The general analytic expression for the Feynman integral of the four-loop vacuum diagram with six propagates for type B in Fig. 3 can be written as

$$\begin{aligned}
U_{6B} &= \left(\Lambda_{\text{RE}}^2 \right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2)[(q_1 + q_3 + q_4)^2 - m_2^2]} \\
&\quad \times \frac{1}{(q_2^2 - m_3^2)[(q_2 + q_3 + q_4)^2 - m_4^2](q_3^2 - m_5^2)(q_4^2 - m_6^2)}.
\end{aligned} \tag{43}$$




FIG. 3: Four-loop vacuum diagram with six propagators for type B, which m_i denotes the mass of the i -th particle and q_j denotes the momentum.

Integrating out \mathbf{q} , the Mellin-Barnes representation of the Feynman integral of the four-loop vacuum diagram with six propagates for type B can be written as

$$\begin{aligned}
U_{6B} = & \frac{m_6^4}{(2\pi i)^5 (4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_6^2} \right)^{8-2D} \int_{-i\infty}^{+i\infty} ds \left[\prod_{i=1}^5 \left(\frac{m_i^2}{m_6^2} \right)^{s_i} \Gamma(-s_i) \right] \\
& \times \left[\prod_{i=1}^5 \Gamma\left(\frac{D}{2} - 1 - s_i\right) \right] \Gamma\left(5 - \frac{3D}{2} + \sum_{i=1}^5 s_i\right) \Gamma\left(6 - 2D + \sum_{i=1}^5 s_i\right) \\
& \times \frac{\Gamma\left(\frac{3D}{2} - 4 - \sum_{i=1}^4 s_i\right) \Gamma\left(2 - \frac{D}{2} + s_1 + s_2\right) \Gamma\left(2 - \frac{D}{2} + s_3 + s_4\right)}{\Gamma\left(4 - D + \sum_{i=1}^4 s_i\right) \Gamma\left(D - 2 - s_1 - s_2\right) \Gamma\left(D - 2 - s_3 - s_4\right)}, \quad (44)
\end{aligned}$$

where $\mathbf{s} = (s_1, \dots, s_5)$.

Taking the residue of the pole of $\Gamma(-s_i)$, ($i = 1, \dots, 5$), one can derive one linear independent term of the integral:

$$U_{6B} \ni \frac{m_6^4}{(4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_6^2} \right)^{8-2D} \frac{\pi^4 \sin^2 \pi D}{\sin^5 \frac{\pi D}{2} \sin \frac{3\pi D}{2}} T_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (45)$$

with

$$T_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) = \sum_{\mathbf{n}=0}^{\infty} A_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}, \quad (46)$$

where $\mathbf{n} = (n_1, \dots, n_5)$, $\mathbf{x} = (x_1, \dots, x_5)$, $x_i = \frac{m_i^2}{m_6^2}$, $\mathbf{a} = (a_1, \dots, a_6)$ and $\mathbf{b} = (b_1, \dots, b_7)$ with

$$\begin{aligned}
a_1 &= 5 - \frac{3D}{2}, \quad a_2 = 6 - 2D, \quad a_3 = a_5 = 2 - \frac{D}{2}, \quad a_4 = a_6 = 3 - D, \\
b_1 &= b_2 = b_3 = b_4 = b_5 = 2 - \frac{D}{2}, \quad b_6 = 4 - D, \quad b_7 = 5 - \frac{3D}{2}, \quad (47)
\end{aligned}$$

and the coefficient A_n is

$$A_n = \frac{\Gamma(a_1 + \sum_{i=1}^5 n_i) \Gamma(a_2 + \sum_{i=1}^5 n_i) \Gamma(a_3 + \sum_{i=1}^2 n_i) \Gamma(a_4 + \sum_{i=1}^2 n_i) \Gamma(a_5 + \sum_{i=3}^4 n_i) \Gamma(a_6 + \sum_{i=3}^4 n_i)}{\left[\prod_{i=1}^5 n_i! \Gamma(b_i + n_i) \right] \Gamma(b_6 + \sum_{i=1}^4 n_i) \Gamma(b_7 + \sum_{i=1}^4 n_i)}. \quad (48)$$

In order to proceed with our analysis, we define the auxiliary function

$$\Phi_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = \mathbf{u}^{\mathbf{a}} \mathbf{v}^{\mathbf{b} - \mathbf{e}_7} T_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (49)$$

with the intermediate variables $\mathbf{u} = (u_1, \dots, u_6) = (1, 1, 1, 1, 1, 1)$, $\mathbf{v} = (v_1, \dots, v_7)$, $\mathbf{v} = \mathbf{e}_7 = (1, 1, 1, 1, 1, 1, 1)$. Then one can obtain

$$\begin{aligned} \vartheta_{u_j} \Phi_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= a_j \Phi_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (j = 1, \dots, 6), \\ \vartheta_{v_k} \Phi_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= (b_k - 1) \Phi_{6B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (k = 1, \dots, 7). \end{aligned} \quad (50)$$

Through the transformation

$$\begin{aligned} z_j &= \frac{1}{u_j}, \quad (j = 1, \dots, 6), \quad z_{6+k} = v_k, \quad (k = 1, \dots, 7), \\ z_{14} &= \frac{x_1}{u_1 u_2 u_3 u_4 v_1 v_6 v_7}, \quad z_{15} = \frac{x_2}{u_1 u_2 u_3 u_4 v_2 v_6 v_7}, \\ z_{16} &= \frac{x_3}{u_1 u_2 u_5 u_6 v_3 v_6 v_7}, \quad z_{17} = \frac{x_4}{u_1 u_2 u_5 u_6 v_4 v_6 v_7}, \quad z_{18} = \frac{x_5}{u_1 u_2 v_5}, \end{aligned} \quad (51)$$

one derives the GKZ hypergeometric system for the four-loop vacuum diagram with six propagates for type B:

$$\mathbf{A}_{6B} \cdot \vec{\vartheta}_{6B} \Phi_{6B} = \mathbf{B}_{6B} \Phi_{6B}, \quad (52)$$

where

$$\begin{aligned} \mathbf{A}_{6B} &= \begin{pmatrix} \mathbf{I}_{13 \times 13} & \mathbf{A}_{\mathbf{x}6B} \end{pmatrix}, \\ \mathbf{A}_{\mathbf{x}6B}^T &= \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & -1 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}, \\ \vec{\vartheta}_{6B}^T &= (\vartheta_{z_1}, \dots, \vartheta_{z_{18}}), \\ \mathbf{B}_{6B}^T &= (-a_1, \dots, -a_6, b_1 - 1, \dots, b_7 - 1). \end{aligned} \quad (53)$$

Here, $\mathbf{I}_{13 \times 13}$ is a 13×13 unit matrix.

Correspondingly the dual matrix $\tilde{\mathbf{A}}_{6\mathbf{B}}$ of $\mathbf{A}_{6\mathbf{B}}$ is

$$\tilde{\mathbf{A}}_{6\mathbf{B}} = \begin{pmatrix} -\mathbf{A}_{\mathbf{X}6\mathbf{B}}^T & \mathbf{I}_{5 \times 5} \end{pmatrix}, \quad (54)$$

where $\mathbf{I}_{5 \times 5}$ is a 5×5 unit matrix. The row vectors of the matrix $\tilde{\mathbf{A}}_{6\mathbf{B}}$ induce the integer sublattice \mathbf{B} which can be used to construct the formal solutions in hypergeometric series.

Defining the combined variables

$$\begin{aligned} y_1 &= \frac{z_7 z_{12} z_{13} z_{14}}{z_1 z_2 z_3 z_4}, & y_2 &= \frac{z_8 z_{12} z_{13} z_{15}}{z_1 z_2 z_3 z_4}, \\ y_3 &= \frac{z_9 z_{12} z_{13} z_{16}}{z_1 z_2 z_5 z_6}, & y_4 &= \frac{z_{10} z_{12} z_{13} z_{17}}{z_1 z_2 z_5 z_6}, & y_5 &= \frac{z_{11} z_{18}}{z_1 z_2}, \end{aligned} \quad (55)$$

we write the solutions satisfying Eq. (52) as

$$\Phi_{6B}(\mathbf{z}) = \left(\prod_{i=1}^{18} z_i^{\alpha_i} \right) \varphi_{6B}(\mathbf{y}), \quad (56)$$

where $\mathbf{y} = (y_1, \dots, y_5)$, $\vec{\alpha}^T = (\alpha_1, \dots, \alpha_{18})$ denotes a sequence of complex number such that

$$\mathbf{A}_{6\mathbf{B}} \cdot \vec{\alpha} = \mathbf{B}_{6\mathbf{B}}. \quad (57)$$

IV. GKZ HYPERGEOMETRIC SYSTEM OF THE FOUR-LOOP VACUUM INTEGRALS WITH SEVEN PROPAGATES

A. Four-loop vacuum diagram with seven propagates for type A

The general analytic expression for the Feynman integral of the four-loop vacuum diagram with seven propagates for type A in Fig. 4 can be written as

$$\begin{aligned} U_{7A} &= \left(\Lambda_{\text{RE}}^2 \right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2)[(q_1 + q_3 + q_4)^2 - m_2^2](q_2^2 - m_3^2)} \\ &\quad \times \frac{1}{[(q_2 + q_3 + q_4)^2 - m_4^2][(q_3 + q_4)^2 - m_5^2](q_3^2 - m_6^2)(q_4^2 - m_7^2)}. \end{aligned} \quad (58)$$

Integrating out \mathbf{q} , the Mellin-Barnes representation of the Feynman integral of the four-loop vacuum diagram with seven propagates for type A can be written as

$$U_{7A} = \frac{-m_7^2}{(2\pi i)^6 (4\pi)^8} \left(\frac{4\pi \Lambda_{\text{RE}}^2}{m_7^2} \right)^{8-2D} \int_{-i\infty}^{+i\infty} ds \left[\prod_{i=1}^6 \left(\frac{m_i^2}{m_7^2} \right)^{s_i} \Gamma(-s_i) \right]$$




FIG. 4: Four-loop vacuum diagram with seven propagators for type A, which m_i denotes the mass of the i -th particle and q_j denotes the momentum.

$$\begin{aligned}
& \times \left[\prod_{i=1}^4 \Gamma\left(\frac{D}{2} - 1 - s_i\right) \right] \Gamma\left(\frac{D}{2} - 1 - s_6\right) \Gamma\left(6 - \frac{3D}{2} + \sum_{i=1}^6 s_i\right) \Gamma\left(7 - 2D + \sum_{i=1}^6 s_i\right) \\
& \times \frac{\Gamma\left(\frac{3D}{2} - 5 - \sum_{i=1}^5 s_i\right) \Gamma\left(2 - \frac{D}{2} + s_1 + s_2\right) \Gamma\left(2 - \frac{D}{2} + s_3 + s_4\right) \Gamma(1 + s_5)}{\Gamma\left(5 - D + \sum_{i=1}^5 s_i\right) \Gamma(D - 2 - s_1 - s_2) \Gamma(D - 2 - s_3 - s_4)}, \quad (59)
\end{aligned}$$

where $\mathbf{s} = (s_1, \dots, s_6)$.

Taking the residue of the pole of $\Gamma(-s_i)$, ($i = 1, \dots, 6$), one can derive one linear independent term of the integral:

$$U_{7A} \ni \frac{-m_7^2}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_7^2} \right)^{8-2D} \frac{\pi^4 \sin^2 \pi D}{\sin^5 \frac{\pi D}{2} \sin \frac{3\pi D}{2}} T_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (60)$$

with

$$T_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) = \sum_{\mathbf{n}=0}^{\infty} A_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}, \quad (61)$$

where $\mathbf{n} = (n_1, \dots, n_6)$, $\mathbf{x} = (x_1, \dots, x_6)$, $x_i = \frac{m_i^2}{m_7^2}$, $\mathbf{a} = (a_1, \dots, a_7)$ and $\mathbf{b} = (b_1, \dots, b_7)$ with

$$\begin{aligned}
a_1 &= 6 - \frac{3D}{2}, \quad a_2 = 7 - 2D, \quad a_3 = a_5 = 2 - \frac{D}{2}, \quad a_4 = a_6 = 3 - D, \quad a_7 = 1, \\
b_1 &= b_2 = b_3 = b_4 = b_5 = 2 - \frac{D}{2}, \quad b_6 = 5 - D, \quad b_7 = 6 - \frac{3D}{2}, \quad (62)
\end{aligned}$$

and the coefficient $A_{\mathbf{n}}$ is

$$A_{\mathbf{n}} = \Gamma\left(a_1 + \sum_{i=1}^6 n_i\right) \Gamma\left(a_2 + \sum_{i=1}^6 n_i\right) \Gamma\left(a_3 + \sum_{i=1}^2 n_i\right)$$

$$\times \frac{\Gamma(a_4 + \sum_{i=1}^2 n_i) \Gamma(a_5 + \sum_{i=3}^4 n_i) \Gamma(a_6 + \sum_{i=3}^4 n_i) \Gamma(a_7 + n_5)}{\left[\prod_{i=1}^6 n_i! \right] \left[\prod_{i=1}^4 \Gamma(b_i + n_i) \right] \Gamma(b_5 + n_6) \Gamma(b_6 + \sum_{i=1}^5 n_i) \Gamma(b_7 + \sum_{i=1}^5 n_i)} . \quad (63)$$

In order to proceed with our analysis, we define the auxiliary function

$$\Phi_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = \mathbf{u}^{\mathbf{a}} \mathbf{v}^{\mathbf{b} - \mathbf{e}_7} T_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) , \quad (64)$$

with the intermediate variables $\mathbf{u} = (u_1, \dots, u_7)$, $\mathbf{v} = (v_1, \dots, v_7)$, $\mathbf{u} = \mathbf{v} = \mathbf{e}_7 = (1, 1, 1, 1, 1, 1, 1)$. Then one can obtain

$$\begin{aligned} \vartheta_{u_j} \Phi_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= a_j \Phi_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) , \quad (j = 1, \dots, 7) , \\ \vartheta_{v_k} \Phi_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) &= (b_k - 1) \Phi_{7A}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) , \quad (k = 1, \dots, 7) . \end{aligned} \quad (65)$$

Through the transformation

$$\begin{aligned} z_j &= \frac{1}{u_j}, \quad (j = 1, \dots, 7), \quad z_{7+k} = v_k, \quad (k = 1, \dots, 7), \\ z_{15} &= \frac{x_1}{u_1 u_2 u_3 u_4 v_1 v_6 v_7}, \quad z_{16} = \frac{x_2}{u_1 u_2 u_3 u_4 v_2 v_6 v_7}, \\ z_{17} &= \frac{x_3}{u_1 u_2 u_5 u_6 v_3 v_6 v_7}, \quad z_{18} = \frac{x_4}{u_1 u_2 u_5 u_6 v_4 v_6 v_7}, \\ z_{19} &= \frac{x_5}{u_1 u_2 u_7 v_6 v_7}, \quad z_{20} = \frac{x_6}{u_1 u_2 v_5}, \end{aligned} \quad (66)$$

one derives the GKZ hypergeometric system for the four-loop vacuum diagram with seven propagates for type A:

$$\mathbf{A}_{7A} \cdot \vec{\vartheta}_{7A} \Phi_{7A} = \mathbf{B}_{7A} \Phi_{7A} , \quad (67)$$

where

$$\begin{aligned} \mathbf{A}_{7A} &= \begin{pmatrix} \mathbf{I}_{14 \times 14} & \mathbf{A}_{\mathbf{x}7A} \end{pmatrix} , \\ \mathbf{A}_{\mathbf{x}7A}^T &= \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix} , \\ \vec{\vartheta}_{7A}^T &= (\vartheta_{z_1}, \dots, \vartheta_{z_{20}}) , \\ \mathbf{B}_{7A}^T &= (-a_1, \dots, -a_7, b_1 - 1, \dots, b_7 - 1) . \end{aligned} \quad (68)$$




FIG. 5: Four-loop vacuum diagram with seven propagators for type B, which m_i denotes the mass of the i -th particle and q_j denotes the momentum.

Here, $\mathbf{I}_{14 \times 14}$ is a 14×14 unit matrix.

Correspondingly the dual matrix $\tilde{\mathbf{A}}_{7\mathbf{A}}$ of $\mathbf{A}_{7\mathbf{A}}$ is

$$\tilde{\mathbf{A}}_{7\mathbf{A}} = \begin{pmatrix} -\mathbf{A}_{7\mathbf{A}}^T & \mathbf{I}_{6 \times 6} \end{pmatrix}, \quad (69)$$

where $\mathbf{I}_{6 \times 6}$ is a 6×6 unit matrix. The row vectors of the matrix $\tilde{\mathbf{A}}_{7\mathbf{A}}$ induce the integer sublattice \mathbf{B} which can be used to construct the formal solutions in hypergeometric series.

Defining the combined variables

$$\begin{aligned} y_1 &= \frac{z_8 z_{13} z_{14} z_{15}}{z_1 z_2 z_3 z_4}, & y_2 &= \frac{z_9 z_{13} z_{14} z_{16}}{z_1 z_2 z_3 z_4}, & y_3 &= \frac{z_{10} z_{13} z_{14} z_{17}}{z_1 z_2 z_5 z_6}, \\ y_4 &= \frac{z_{11} z_{13} z_{14} z_{18}}{z_1 z_2 z_5 z_6}, & y_5 &= \frac{z_{13} z_{14} z_{19}}{z_1 z_2 z_7}, & y_6 &= \frac{z_{12} z_{20}}{z_1 z_2}, \end{aligned} \quad (70)$$

we write the solutions satisfying Eq. (67) as

$$\Phi_{7\mathbf{A}}(\mathbf{z}) = \left(\prod_{i=1}^{20} z_i^{\alpha_i} \right) \varphi_{7\mathbf{A}}(\mathbf{y}), \quad (71)$$

where $\mathbf{y} = (y_1, \dots, y_6)$, and $\vec{\alpha}^T = (\alpha_1, \alpha_2, \dots, \alpha_{20})$ denotes a sequence of complex number such that

$$\mathbf{A}_{7\mathbf{A}} \cdot \vec{\alpha} = \mathbf{B}_{7\mathbf{A}}. \quad (72)$$

B. Four-loop vacuum diagram with seven propagates for type B

The general analytic expression for the Feynman integral of the four-loop vacuum diagram with seven propagates for type B in Fig. 5 is

$$U_{7B} = \left(\Lambda_{\text{RE}}^2\right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2)[(q_1 + q_2)^2 - m_2^2](q_2^2 - m_3^2)} \\ \times \frac{1}{[(q_2 + q_3 + q_4)^2 - m_4^2][(q_3 + q_4)^2 - m_5^2](q_3^2 - m_6^2)(q_4^2 - m_7^2)}. \quad (73)$$

Integrating out \mathbf{q} , the Mellin-Barnes representation of the Feynman integral of the four-loop vacuum diagram with seven propagates for type B can be written as

$$U_{7B} = \frac{-m_7^2}{(2\pi i)^6 (4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_7^2}\right)^{8-2D} \int_{-i\infty}^{+i\infty} ds \left[\prod_{i=1}^6 \left(\frac{m_i^2}{m_7^2}\right)^{s_i} \Gamma(-s_i) \right] \left[\prod_{i=1}^2 \Gamma\left(\frac{D}{2} - 1 - s_i\right) \right] \\ \times \Gamma\left(\frac{D}{2} - 1 - s_4\right) \Gamma\left(\frac{D}{2} - 1 - s_6\right) \Gamma\left(6 - \frac{3D}{2} + \sum_{i=1}^6 s_i\right) \Gamma\left(7 - 2D + \sum_{i=1}^6 s_i\right) \Gamma\left(4 - D + \sum_{i=1}^4 s_i\right) \\ \times \frac{\Gamma\left(\frac{3D}{2} - 5 - \sum_{i=1}^5 s_i\right) \Gamma\left(D - 3 - \sum_{i=1}^3 s_i\right) \Gamma\left(2 - \frac{D}{2} + s_1 + s_2\right) \Gamma(1 + s_3) \Gamma(1 + s_5)}{\Gamma(D - 2 - s_1 - s_2) \Gamma\left(\frac{3D}{2} - 4 - \sum_{i=1}^4 s_i\right) \Gamma\left(3 - \frac{D}{2} + \sum_{i=1}^3 s_i\right) \Gamma(5 - D + \sum_{i=1}^5 s_i)}, \quad (74)$$

where $\mathbf{s} = (s_1, \dots, s_6)$.

Taking the residue of the pole of $\Gamma(-s_i)$, ($i = 1, \dots, 6$), one can derive one linear independent term of the integral:

$$U_{7B} \ni \frac{m_7^2}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_7^2}\right)^{8-2D} \frac{\pi^4}{\sin^4 \frac{\pi D}{2}} T_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (75)$$

with

$$T_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}) = \sum_{\mathbf{n}=0}^{\infty} A_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}, \quad (76)$$

where $\mathbf{n} = (n_1, \dots, n_6)$, $\mathbf{x} = (x_1, \dots, x_6)$, $x_i = \frac{m_i^2}{m_7^2}$, $\mathbf{a} = (a_1, \dots, a_8)$ and $\mathbf{b} = (b_1, \dots, b_8)$ with

$$a_1 = 6 - \frac{3D}{2}, \quad a_2 = 7 - 2D, \quad a_3 = 5 - \frac{3D}{2}, \quad a_4 = 4 - D, \\ a_5 = 2 - \frac{D}{2}, \quad a_6 = 3 - D, \quad a_7 = a_8 = 1, \quad b_1 = b_2 = b_3 = b_4 = 2 - \frac{D}{2}, \\ b_5 = 3 - \frac{D}{2}, \quad b_6 = 4 - D, \quad b_7 = 5 - D, \quad b_8 = 6 - \frac{3D}{2}, \quad (77)$$

and the coefficient $A_{\mathbf{n}}$ is

$$A_{\mathbf{n}} = \frac{\Gamma(a_1 + \sum_{i=1}^6 n_i) \Gamma(a_2 + \sum_{i=1}^6 n_i) \Gamma(a_3 + \sum_{i=1}^4 n_i) \Gamma(a_4 + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^6 n_i! \right] \Gamma(b_1 + n_1) \Gamma(b_2 + n_2) \Gamma(b_3 + n_4) \Gamma(b_4 + n_6)} \\ \times \frac{\Gamma(a_5 + \sum_{i=1}^2 n_i) \Gamma(a_6 + \sum_{i=1}^2 n_i) \Gamma(a_7 + n_3) \Gamma(a_8 + n_5)}{\Gamma(b_5 + \sum_{i=1}^3 n_i) \Gamma(b_6 + \sum_{i=1}^3 n_i) \Gamma(b_7 + \sum_{i=1}^5 n_i) \Gamma(b_8 + \sum_{i=1}^5 n_i)}. \quad (78)$$

We also define the auxiliary function

$$\Phi_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = \mathbf{u}^{\mathbf{a}} \mathbf{v}^{\mathbf{b} - \mathbf{e}_8} T_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}), \quad (79)$$

with the intermediate variables $\mathbf{u} = (u_1, \dots, u_8)$, $\mathbf{v} = (v_1, \dots, v_8)$, $\mathbf{u} = \mathbf{v} = \mathbf{e}_8 = (1, 1, 1, 1, 1, 1, 1, 1)$. Then one can obtain

$$\vartheta_{u_j} \Phi_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = a_j \Phi_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (j = 1, \dots, 8), \\ \vartheta_{v_k} \Phi_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}) = (b_k - 1) \Phi_{7B}(\mathbf{a}, \mathbf{b} \mid \mathbf{x}, \mathbf{u}, \mathbf{v}), \quad (k = 1, \dots, 8). \quad (80)$$

Through the transformation

$$z_j = \frac{1}{u_j}, \quad (j = 1, \dots, 8), \quad z_{8+k} = v_k, \quad (k = 1, \dots, 8), \\ z_{17} = \frac{x_1}{u_1 u_2 u_3 u_4 u_5 u_6 v_1 v_5 v_6 v_7 v_8}, \quad z_{18} = \frac{x_2}{u_1 u_2 u_3 u_4 u_5 u_6 v_2 v_5 v_6 v_7 v_8}, \\ z_{19} = \frac{x_3}{u_1 u_2 u_3 u_4 u_7 v_5 v_6 v_7 v_8}, \quad z_{20} = \frac{x_4}{u_1 u_2 u_3 u_4 v_3 v_7 v_8}, \\ z_{21} = \frac{x_5}{u_1 u_2 u_8 v_7 v_8}, \quad z_{22} = \frac{x_6}{u_1 u_2 v_4}, \quad (81)$$

one derives the GKZ hypergeometric system for the four-loop vacuum diagram with seven propagates for type B:

$$\mathbf{A}_{7B} \cdot \vec{\vartheta}_{7B} \Phi_{7B} = \mathbf{B}_{7B} \Phi_{7B}, \quad (82)$$

where

$$\mathbf{A}_{7B} = \begin{pmatrix} \mathbf{I}_{16 \times 16} & \mathbf{A}_{\mathbf{x}7B} \end{pmatrix},$$

$$\mathbf{A}_{\mathbf{x7B}}^T = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & -1 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\vec{\vartheta}_{7B}^T = (\vartheta_{z_1}, \dots, \vartheta_{z_{22}}),$$

$$\mathbf{B}_{7B}^T = (-a_1, \dots, -a_8, b_1 - 1, \dots, b_8 - 1). \quad (83)$$

Here, $\mathbf{I}_{16 \times 16}$ is a 16×16 unit matrix.

Correspondingly the dual matrix $\tilde{\mathbf{A}}_{7B}$ of \mathbf{A}_{7B} is

$$\tilde{\mathbf{A}}_{7B} = \begin{pmatrix} -\mathbf{A}_{\mathbf{x7B}}^T & \mathbf{I}_{6 \times 6} \end{pmatrix}, \quad (84)$$

where $\mathbf{I}_{6 \times 6}$ is a 6×6 unit matrix. The row vectors of the matrix $\tilde{\mathbf{A}}_{7B}$ also induce the integer sublattice \mathbf{B} which can be used to construct the formal solutions in hypergeometric series.

Defining the combined variables

$$\begin{aligned} y_1 &= \frac{z_9 z_{13} z_{14} z_{15} z_{16} z_{17}}{z_1 z_2 z_3 z_4 z_5 z_6}, & y_2 &= \frac{z_{10} z_{13} z_{14} z_{15} z_{16} z_{18}}{z_1 z_2 z_3 z_4 z_5 z_6}, \\ y_3 &= \frac{z_{13} z_{14} z_{15} z_{16} z_{19}}{z_1 z_2 z_3 z_4 z_7}, & y_4 &= \frac{z_{11} z_{15} z_{16} z_{20}}{z_1 z_2 z_3 z_4}, \\ y_5 &= \frac{z_{15} z_{16} z_{21}}{z_1 z_2 z_8}, & y_6 &= \frac{z_{12} z_{22}}{z_1 z_2}, \end{aligned} \quad (85)$$

we write the solutions satisfying Eq. (82) as

$$\Phi_{7B}(\mathbf{z}) = \left(\prod_{i=1}^{22} z_i^{\alpha_i} \right) \varphi_{7B}(\mathbf{y}), \quad (86)$$

where $\vec{\alpha}^T = (\alpha_1, \dots, \alpha_{22})$ denotes a sequence of complex number such that

$$\mathbf{A}_{7B} \cdot \vec{\alpha} = \mathbf{B}_{7B}. \quad (87)$$

V. THE HYPERGEOMETRIC SERIES SOLUTIONS OF THE FOUR-LOOP VACUUM INTEGRALS

A. The general case of the four-loop vacuum integral with five propagates

In this subsection, we will show the hypergeometric series solutions of the GKZ hypergeometric system of the four-loop vacuum integral with five propagates with arbitrary masses. To construct the hypergeometric series solutions of the GKZ hypergeometric system of the four-loop vacuum integral with five propagates in Eq. (18) is equivalent to choose a set of the linear independent column vectors of the matrix in Eq. (20) which spans the dual space. We denote the submatrix composed of the first, third, fourth and fifth column vectors of the dual matrix of Eq. (20) as $\tilde{\mathbf{A}}_{1345}$, i.e.

$$\tilde{\mathbf{A}}_{1345} = \begin{pmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{pmatrix}. \quad (88)$$

Obviously $\det \tilde{\mathbf{A}}_{1345} = 1 \neq 0$, and

$$\begin{aligned} \mathbf{B}_{1345} &= \tilde{\mathbf{A}}_{1345}^{-1} \cdot \tilde{\mathbf{A}}_5 \\ &= \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & -1 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 & 1 & -1 \end{pmatrix}. \end{aligned} \quad (89)$$

Taking 4 row vectors of the matrix \mathbf{B}_{1345} as the basis of integer lattice, one constructs the GKZ hypergeometric series solutions in parameter space through choosing the sets of column indices $I_i \subset [1, \dots, 10]$ ($i = 1, \dots, 16$) which are consistent with the basis of integer lattice \mathbf{B}_{1345} .

We take the set of column indices $I_1 = [2, 6, \dots, 10]$, i.e. the implement $J_1 = [1, \dots, 10] \setminus I_1 = [1, 3, 4, 5]$. The choice on the set of indices implies the exponent numbers $\alpha_1 = \alpha_3 = \alpha_4 = \alpha_5 = 0$. Through Eq. (24), one can have

$$\alpha_2 = a_1 - a_2, \quad \alpha_6 = \sum_{i=1}^4 b_i - a_1 - 4, \quad \alpha_7 = 1 - b_1,$$

$$\alpha_8 = 1 - b_2, \alpha_9 = 1 - b_3, \alpha_{10} = \sum_{i=1}^3 b_i - a_1 - 3. \quad (90)$$

Combined with Eq. (9), we can have

$$\alpha_2 = \frac{D}{2} - 1, \alpha_6 = -\frac{D}{2}, \alpha_7 = \alpha_8 = \alpha_9 = \frac{D}{2} - 1, \alpha_{10} = -1. \quad (91)$$

According the basis of integer lattice \mathbf{B}_{1345} , the corresponding hypergeometric series solution with quadruple independent variables is written as

$$\begin{aligned} \Phi_{[1345]}^{(1)} &= \prod_{i=1}^{10} z_i^{\alpha_i} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(1)}(\mathbf{n}) \left(\frac{z_1 z_2}{z_6 z_{10}} \right)^{n_1} \left(\frac{z_3 z_7}{z_6 z_{10}} \right)^{n_2} \left(\frac{z_4 z_8}{z_6 z_{10}} \right)^{n_3} \left(\frac{z_5 z_9}{z_6 z_{10}} \right)^{n_4} \\ &= \prod_{i=1}^{10} z_i^{\alpha_i} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(1)}(\mathbf{n}) \left(\frac{1}{y_4} \right)^{n_1} \left(\frac{y_1}{y_4} \right)^{n_2} \left(\frac{y_2}{y_4} \right)^{n_3} \left(\frac{y_3}{y_4} \right)^{n_4}, \end{aligned} \quad (92)$$

where the coefficient is

$$\begin{aligned} c_{[1345]}^{(1)}(\mathbf{n}) &= \left\{ \left[\prod_{i=1}^4 n_i! \right] \Gamma(1 + \alpha_2 + n_1) \Gamma(1 + \alpha_6 - \sum_{i=1}^4 n_i) \Gamma(1 + \alpha_7 + n_2) \right. \\ &\quad \left. \times \Gamma(1 + \alpha_8 + n_3) \Gamma(1 + \alpha_9 + n_4) \Gamma(1 + \alpha_{10} - \sum_{i=1}^4 n_i) \right\}^{-1}. \end{aligned} \quad (93)$$

Using the relation in Eq. (6), one can have

$$c_{[1345]}^{(1)}(\mathbf{n}) \propto \frac{\Gamma(-\alpha_6 + \sum_{i=1}^4 n_i) \Gamma(-\alpha_{10} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(1 + \alpha_2 + n_1) \Gamma(1 + \alpha_7 + n_2) \Gamma(1 + \alpha_8 + n_3) \Gamma(1 + \alpha_9 + n_4)}, \quad (94)$$

where we ignore the constant coefficient term $\frac{\sin \pi \alpha_6 \sin \pi \alpha_{10}}{\pi^2}$. And then, through Eq. (91), the corresponding hypergeometric series solution can be written as

$$\begin{aligned} \Phi_{[1345]}^{(1)} &= y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{-1} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(1)}(\mathbf{n}) f_{[1345]}, \\ f_{[1345]} &= \left(\frac{1}{y_4} \right)^{n_1} \left(\frac{y_1}{y_4} \right)^{n_2} \left(\frac{y_2}{y_4} \right)^{n_3} \left(\frac{y_3}{y_4} \right)^{n_4}, \end{aligned} \quad (95)$$

with the coefficient is

$$c_{[1345]}^{(1)}(\mathbf{n}) = \frac{\Gamma(\frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(1 + \sum_{i=1}^4 n_i)}{\prod_{i=1}^4 n_i! \Gamma(\frac{D}{2} + n_i)}. \quad (96)$$

Here, the convergent region of the hypergeometric function $\Phi_{[1345]}^{(1)}$ in Eq. (95) is

$$\Xi_{[1345]} = \{(y_1, y_2, y_3, y_4) | 1 < |y_4|, |y_1| < |y_4|, |y_2| < |y_4|, |y_3| < |y_4|\}, \quad (97)$$

which shows that $\Phi_{[1345]}^{(1)}$ is in neighborhood of regular singularity ∞ .

According the basis of integer lattice \mathbf{B}_{1345} , we can also obtain other fifteen hypergeometric solutions, which the expressions are collected in Appendix B. The sixteen hypergeometric series solutions $\Phi_{[1345]}^{(i)}$ whose convergent region is $\Xi_{[1345]}$ can constitute a fundamental solution system. The combination coefficients are determined by the value of the scalar integral of an ordinary point or some regular singularities.

Multiplying one of the row vectors of the matrix \mathbf{B}_{1345} by -1, the induced integer matrix can also be chosen as a basis of the integer lattice space of certain hypergeometric series. Taking 4 row vectors of the following matrix as the basis of integer lattice,

$$\begin{aligned} \mathbf{B}_{1345} &= \text{diag}(-1, 1, 1, 1) \cdot \mathbf{B}_{1345} \\ &= \begin{pmatrix} -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & -1 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 & 1 & -1 \end{pmatrix}, \end{aligned} \quad (98)$$

one obtains sixteen hypergeometric series solutions $\Phi_{[\bar{1}345]}^{(i)}$ ($i = 1, \dots, 16$) similarly, which the expressions are collected in Appendix C. The convergent region of the hypergeometric functions $\Phi_{[\bar{1}345]}^{(i)}$ is

$$\Xi_{[\bar{1}345]} = \{(y_1, y_2, y_3, y_4) \mid |y_1| < 1, |y_2| < 1, |y_3| < 1, |y_4| < 1\}, \quad (99)$$

which shows that $\Phi_{[\bar{1}345]}^{(i)}$ are in neighborhood of regular singularity 0 and can constitute a fundamental solution system.

Taking 4 row vectors of the following matrix as the basis of integer lattice,

$$\begin{aligned} \mathbf{B}_{1\bar{3}45} &= \text{diag}(1, -1, 1, 1) \cdot \mathbf{B}_{1345} \\ &= \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 & 1 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 & 1 & -1 \end{pmatrix}, \end{aligned} \quad (100)$$

one also obtains sixteen hypergeometric series solutions $\Phi_{[1\bar{3}45]}^{(i)}$ ($i = 1, \dots, 16$):

$$\Phi_{[1\bar{3}45]}^{(i)} = \Phi_{[1345]}^{(i)} (y_4 \leftrightarrow y_1), \quad (101)$$

which the expressions can be obtained by interchanging between y_4 and y_1 in $\Phi_{[1345]}^{(i)}$. The convergent region of the hypergeometric functions $\Phi_{[1345]}^{(i)}$ is

$$\Xi_{[1\bar{3}45]} = \{(y_1, y_2, y_3, y_4) | 1 < |y_1|, |y_4| < |y_1|, |y_2| < |y_1|, |y_3| < |y_1|\} , \quad (102)$$

which shows that $\Phi_{[1345]}^{(i)}$ are in neighborhood of regular singularity ∞ and can constitute a fundamental solution system.

Taking 4 row vectors of the following matrix as the basis of integer lattice,

$$\begin{aligned} \mathbf{B}_{134\bar{5}} &= \text{diag}(1, 1, -1, 1) \cdot \mathbf{B}_{1345} \\ &= \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & -1 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 & 1 & -1 \end{pmatrix} , \end{aligned} \quad (103)$$

one obtains sixteen hypergeometric series solutions $\Phi_{[1345]}^{(i)}$ ($i = 1, \dots, 16$):

$$\Phi_{[1345]}^{(i)} = \Phi_{[1345]}^{(i)}(y_4 \leftrightarrow y_2) , \quad (104)$$

which the expressions can be obtained by interchanging between y_4 and y_2 in $\Phi_{[1345]}^{(i)}$. The convergent region of the hypergeometric functions $\Phi_{[1345]}^{(i)}$ is

$$\Xi_{[134\bar{5}]} = \{(y_1, y_2, y_3, y_4) | 1 < |y_2|, |y_1| < |y_2|, |y_4| < |y_2|, |y_3| < |y_2|\} , \quad (105)$$

which shows that $\Phi_{[1345]}^{(i)}$ are in neighborhood of regular singularity ∞ and can constitute a fundamental solution system.

Taking 4 row vectors of the following matrix as the basis of integer lattice,

$$\begin{aligned} \mathbf{B}_{134\bar{5}} &= \text{diag}(1, 1, 1, -1) \cdot \mathbf{B}_{1345} \\ &= \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & -1 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 & -1 & 1 \end{pmatrix} , \end{aligned} \quad (106)$$

one obtains sixteen hypergeometric series solutions $\Phi_{[1345]}^{(i)}$ ($i = 1, \dots, 16$):

$$\Phi_{[134\bar{5}]}^{(i)} = \Phi_{[1345]}^{(i)}(y_4 \leftrightarrow y_3) , \quad (107)$$

which the expressions can be obtained by interchanging between y_4 and y_3 in $\Phi_{[1345]}^{(i)}$. The convergent region of the hypergeometric functions $\Phi_{[134\bar{5}]}^{(i)}$ is

$$\Xi_{[134\bar{5}]} = \{(y_1, y_2, y_3, y_4) \mid 1 < |y_3|, |y_1| < |y_3|, |y_2| < |y_3|, |y_4| < |y_3|\} , \quad (108)$$

which shows that $\Phi_{[134\bar{5}]}^{(i)}$ are in neighborhood of regular singularity ∞ and can constitute a fundamental solution system.

B. The special case of the four-loop vacuum integral with five propagates

In order to elucidate how to obtain the analytical expression clearly, we assume the two nonzero virtual mass for the four-loop vacuum integral with five propagates. The corresponding integral of the special case for the four-loop vacuum diagram can be expressed as a linear combination of those corresponding functionally independent Gauss functions.

Through Sec. II, the GKZ hypergeometric system in this special case ($m_1 \neq 0$, $m_5 \neq 0$, $m_2 = m_3 = m_4 = 0$) can be simplified as

$$\mathbf{A}_{51} \cdot \vec{\vartheta}_{51} \Phi_{51} = \mathbf{B}_5 \Phi_{51} , \quad (109)$$

where the vector of Euler operators is defined as

$$\vec{\vartheta}_{51}^T = (\vartheta_{z_1}, \dots, \vartheta_{z_7}) , \quad (110)$$

and the matrix \mathbf{A}_{51} is obtained through deleting the 8th, 9th, and 10th columns of the matrix \mathbf{A}_5 :

$$\mathbf{A}_{51} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix} . \quad (111)$$

Through Eq. (109), one can have the relations

$$\alpha_1 + \alpha_7 = -a_1 , \quad \alpha_2 + \alpha_7 = -a_2 , \quad \alpha_3 - \alpha_7 = b_1 - 1 , \quad (112)$$

with $a_1 = 4 - \frac{3D}{2}$, $a_2 = 5 - 2D$, $b_1 = 2 - \frac{D}{2}$, and the other α_i are zero. The dual matrix $\tilde{\mathbf{A}}_{51}$ of \mathbf{A}_{51} is

$$\tilde{\mathbf{A}}_{51} = \begin{pmatrix} -1 & -1 & 1 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (113)$$

The integer sublattice \mathbf{B}_{51} is determined by the dual matrix $\tilde{\mathbf{A}}_{51}$ with $\mathbf{B}_{51} = \tilde{\mathbf{A}}_{51}$. The integer sublattice \mathbf{B}_{51} implies that the system of fundamental solutions is composed by two linear independent hypergeometric functions. We take the set of column indices $I_1 = [1, \dots, 6]$, which implies the exponent numbers $\alpha_7 = 0$. And then, the corresponding hypergeometric series solution can be written as

$$\Phi_{[51]}^{(1)}(y_1) = {}_2F_1 \left(\begin{matrix} 4 - \frac{3D}{2}, & 5 - 2D \\ & 2 - \frac{D}{2} \end{matrix} \middle| y_1 \right), \quad (114)$$

with $y_1 = x_1 = m_1^2/m_5^2$, and ${}_2F_1$ is Gauss function:

$${}_2F_1 \left(\begin{matrix} a, & b \\ & c \end{matrix} \middle| x \right) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n! (c)_n} x^n, \quad (115)$$

with $(a)_n = \Gamma(a+n)/\Gamma(a)$. We also can take the set of column indices $I_2 = [1, 2, 4, \dots, 7]$, which implies the exponent numbers $\alpha_3 = 0$. And then, the another hypergeometric series solution can be written as

$$\Phi_{[51]}^{(2)}(y_1) = (y_1)^{D/2-1} {}_2F_1 \left(\begin{matrix} 3 - D, & 4 - \frac{3D}{2} \\ & \frac{D}{2} \end{matrix} \middle| y_1 \right). \quad (116)$$

Here, the convergent region of the hypergeometric functions $\Phi_{[51]}^{(1,2)}(y_1)$ is $|y_1| < 1$. In the region $|y_1| < 1$, the integral correspondingly is a linear combination of two fundamental solutions:

$$\Phi_{51}(y_1) = C_{[51]}^{(1)} \Phi_{[51]}^{(1)}(y_1) + C_{[51]}^{(2)} \Phi_{[51]}^{(2)}(y_1). \quad (117)$$

Multiplying one of the row vectors of the integer matrix \mathbf{B}_{51} by -1, the induced integer matrix also can be chosen as a basis of the integer lattice space of certain hypergeometric series. And the corresponding system of fundamental solutions for is similarly composed by two Gauss functions:

$$\begin{aligned} \Phi_{[51]}^{(3)}(y_1) &= (y_1)^{\frac{3D}{2}-4} {}_2F_1 \left(\begin{matrix} 4 - \frac{3D}{2}, & 3 - D \\ & \frac{D}{2} \end{matrix} \middle| \frac{1}{y_1} \right), \\ \Phi_{[51]}^{(4)}(y_1) &= (y_1)^{2D-5} {}_2F_1 \left(\begin{matrix} 5 - 2D, & 4 - \frac{3D}{2} \\ & 2 - \frac{D}{2} \end{matrix} \middle| \frac{1}{y_1} \right), \end{aligned} \quad (118)$$

which the convergent region is $|y_1| > 1$. Correspondingly the integral in the region $|y_1| > 1$ is a linear combination of two fundamental solutions:

$$\Phi_{51}(y_1) = C_{[51]}^{(3)} \Phi_{[51]}^{(3)}(y_1) + C_{[51]}^{(4)} \Phi_{[51]}^{(4)}(y_1). \quad (119)$$

As $m_1^2 \ll m_5^2$, $m_2 = m_3 = m_4 = 0$,

$$\begin{aligned} I_1 &= \left(\Lambda_{\text{RE}}^2\right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2) q_2^2 q_3^2 (q_1 + q_2 + q_3 + q_4)^2 (q_3^2 - m_5^2)} \\ &= I_{1,0} + \dots, \end{aligned} \quad (120)$$

where

$$\begin{aligned} I_{1,0} &= \left(\Lambda_{\text{RE}}^2\right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{q_1^2 q_2^2 q_3^2 (q_1 + q_2 + q_3 + q_4)^2 (q_3^2 - m_5^2)} \\ &= \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2}\right)^{8-2D} \Gamma(4 - \frac{3D}{2}) \Gamma(5 - 2D). \end{aligned} \quad (121)$$

This result indicates

$$C_{[51]}^{(1)} = \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2}\right)^{8-2D} \Gamma(4 - \frac{3D}{2}) \Gamma(5 - 2D). \quad (122)$$

As $m_1^2 \gg m_5^2$ and $m_2 = m_3 = m_4 = 0$, $I_1 = I_{1,\infty} + \dots$, where

$$\begin{aligned} I_{1,\infty} &= \left(\Lambda_{\text{RE}}^2\right)^{8-2D} \int \frac{d^D \mathbf{q}}{(2\pi)^D} \frac{1}{(q_1^2 - m_1^2) q_2^2 q_3^2 (q_1 + q_2 + q_3 + q_4)^2 q_3^2} \\ &= \frac{-m_1^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_1^2}\right)^{8-2D} \Gamma(4 - \frac{3D}{2}) \Gamma(5 - 2D). \end{aligned} \quad (123)$$

This result indicates

$$C_{[51]}^{(4)} = \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2}\right)^{8-2D} \Gamma(4 - \frac{3D}{2}) \Gamma(5 - 2D). \quad (124)$$

Actually, the Mellin-Barnes representation of the Feynman integral in this case can be obtained as

$$\begin{aligned} U_5 &= \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2}\right)^{8-2D} \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} ds_1 \left(\frac{m_1^2}{m_5^2}\right)^{s_1} \Gamma(-s_1) \\ &\quad \times \Gamma\left(\frac{D}{2} - 1 - s_1\right) \Gamma\left(4 - \frac{3D}{2} + s_1\right) \Gamma(5 - 2D + s_1). \end{aligned} \quad (125)$$

The residue of simple pole of $\Gamma(-s_1)$ provides $C_{[51]}^{(1)} \Phi_{[51]}^{(1)}(y_1)$, that of simple pole of $\Gamma(D/2 - 1 - s_1)$ provides $C_{[51]}^{(2)} \Phi_{[51]}^{(2)}(y_1)$, that of simple pole of $\Gamma(4 - \frac{3D}{2} + s_1)$ provides $C_{[51]}^{(3)} \Phi_{[51]}^{(3)}(y_1)$, and

that of simple pole of $\Gamma(5 - 2D + s_1)$ provides $C_{[51]}^{(4)}\Phi_{[51]}^{(4)}(y_1)$, respectively. And the residue of the simple pole of $\Gamma(D/2 - 1 - s_1)$ and $\Gamma(4 - \frac{3D}{2} + s_1)$ can induce

$$C_{[51]}^{(2)} = C_{[51]}^{(3)} = \frac{-m_5^6}{(4\pi)^8} \left(\frac{4\pi\Lambda_{\text{RE}}^2}{m_5^2} \right)^{8-2D} \Gamma(4 - \frac{3D}{2}) \Gamma(3 - D) \Gamma(1 - \frac{D}{2}). \quad (126)$$

A conclusion for the other two nonzero virtual mass for the four-loop vacuum integral with five propagates, such as $m_2 \neq 0$, $m_5 \neq 0$, $m_1 = m_3 = m_4 = 0$ is analogous to that of $m_1 \neq 0$, $m_5 \neq 0$, $m_2 = m_3 = m_4 = 0$.

C. The four-loop vacuum integrals with six or seven propagates

In our previous works [85, 86], we obtain GKZ hypergeometric systems of some one-loop and two-loop Feynman integrals, which show the algorithm and the obvious hypergeometric series solutions for the one-loop and two-loop Feynman integrals. Recently, the authors of the Refs. [72, 103] also give publicly available computer packages MBConicHulls [72] and FeynGKZ [103] to compute Feynman integrals in terms of hypergeometric functions, which are meaningful to improve computing efficiency. Through the package FeynGKZ [103], they give the examples of some one-loop and two-loop Feynman integrals, which are tested analytically by our previous work [85, 86], as well as numerically using the package FIESTA [109].

Here, we also evaluate the four-loop vacuum integrals with six or seven propagates using FeynGKZ [103], which can be seen in the supplementary material. Note that FeynGKZ can't evaluate the four-loop vacuum integrals with five propagates through Mellin-Barnes representations and Miller's transformation. In the supplementary material, we can see that the GKZ hypergeometric systems of four-loop vacuum integrals with six or seven propagates are in agree with our above results. Using FeynGKZ, we also give some hypergeometric series solutions for the four-loop vacuum integrals with six or seven propagates. The series solutions from GKZ hypergeometric systems for special case with the two nonzero virtual masses and the three nonzero virtual masses are also showed in the supplementary material, and tested numerically using FIESTA [109]. One can see that the computing time using the hypergeometric series solutions is about $\mathcal{O}(10^{-5})$ times that using FIESTA, which evaluate quickly.

Note that, except above five topologies, the four-loop vacuum integrals still have five topologies, which are one topology with seven propagates, two topologies with eight prop-

agates, and two topologies with nine propagates [44]. Due that they have complex mathematical structure, the five topologies of the four-loop vacuum integrals can't obtain the GKZ hypergeometric systems, through Mellin-Barnes representations and Miller's transformation. In next work, we will embed the general four-loop vacuum Feynman integrals into the subvarieties of Grassmannian manifold [87], to explore more possibilities of the general four-loop vacuum Feynman integrals.

VI. CONCLUSIONS

Using Mellin-Barnes representation and Miller's transformation, we derive GKZ hypergeometric systems of the four-loop vacuum integrals with arbitrary masses. The dimension of the GKZ hypergeometric system equals the number of independent dimensionless ratios among the virtual mass squared. In the neighborhoods of origin including infinity, we can obtain analytical hypergeometric series solutions of the four-loop vacuum integrals through GKZ hypergeometric systems. The linear independent hypergeometric series solutions whose convergent regions have non-empty intersection can constitute a fundamental solution system in a proper subset of the whole parameter space. In certain convergent region, the four-loop vacuum integrals can be formulated as a linear combination of the corresponding fundamental solution system. The combination coefficients are determined by the vacuum integral at some ordinary points or regular singularities, or the Mellin-Barnes representation of the vacuum integral.

Here, we obtain the analytical hypergeometric solutions of the four-loop vacuum integrals in the neighborhoods of origin including infinity, using GKZ hypergeometric systems on general manifold. In order to derive the fundamental solution system in neighborhoods of all possible regular singularities, we can embed the vacuum integrals in corresponding Grassmannian manifold [87] through their parametrization. To efficiently derive the fundamental solution system, next we will embed the four-loop vacuum integrals into the subvarieties of Grassmannian using the α -parametric representation.

Acknowledgments

The work has been supported by the National Natural Science Foundation of China (NNSFC) with Grants No. 12075074, No. 12235008, Hebei Natural Science Foundation with Grants No. A2022201017, No. A2023201041, Natural Science Foundation of Guangxi Autonomous Region with Grant No. 2022GXNSFDA035068, and the youth top-notch talent support program of the Hebei Province.

APPENDIX A: THE CALCULATION OF THE INTEGRAL I_q

For the integral I_q , firstly we can integrate out q_1 :

$$\begin{aligned} I_{q_1} &\equiv \int \frac{d^D q_1}{(2\pi)^D} \frac{1}{(q_1^2)^{1+s_1} ((q_1 + q_2 + q_3 + q_4)^2)^{1+s_4}} \\ &= \frac{\Gamma(2 + s_1 + s_4)}{\Gamma(1 + s_1)\Gamma(1 + s_4)} \int_0^1 dx (1-x)^{s_1} x^{s_4} \\ &\quad \times \int \frac{d^D q_1}{(2\pi)^D} \frac{1}{[q_1^2 + x(1-x)(q_2 + q_3 + q_4)^2]^{2+s_1+s_4}}. \end{aligned} \quad (\text{A1})$$

Using the well-known integral

$$\int \frac{d^D q}{(2\pi)^D} \frac{1}{[q^2 + \Delta]^n} = \frac{i(-)^{D/2}\Gamma(n - \frac{D}{2})}{(4\pi)^{D/2}\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n-D/2}, \quad (\text{A2})$$

one can have

$$\begin{aligned} I_{q_1} &= \frac{i(-)^{D/2}\Gamma(2 - \frac{D}{2} + s_1 + s_4)}{(4\pi)^{D/2}\Gamma(1 + s_1)\Gamma(1 + s_4)} \frac{1}{[(q_2 + q_3 + q_4)^2]^{2-\frac{D}{2}+s_1+s_4}} \\ &\quad \times \int_0^1 dx x^{D/2-2-s_1} (1-x)^{D/2-2-s_4}. \end{aligned} \quad (\text{A3})$$

Through Beta function

$$B(m, n) = \int_0^1 dx x^{m-1} (1-x)^{n-1} = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}, \quad (\text{A4})$$

we can have

$$\begin{aligned} I_{q_1} &= \frac{i(-)^{D/2}\Gamma(2 - \frac{D}{2} + s_1 + s_4)\Gamma(\frac{D}{2} - 1 - s_1)\Gamma(\frac{D}{2} - 1 - s_4)}{(4\pi)^{D/2}\Gamma(1 + s_1)\Gamma(1 + s_4)\Gamma(D - 2 - s_1 - s_4)} \\ &\quad \times \frac{1}{[(q_2 + q_3 + q_4)^2]^{2-\frac{D}{2}+s_1+s_4}}. \end{aligned} \quad (\text{A5})$$

Similarly, we can integrate out q_2 :

$$\begin{aligned}
I_{q_2} &\equiv \int \frac{d^D q_2}{(2\pi)^D} \frac{1}{(q_2^2)^{1+s_2} [(q_2 + q_3 + q_4)^2]^{2-\frac{D}{2}+s_1+s_4}} \\
&= \frac{i(-)^{D/2} \Gamma(3-D+s_1+s_2+s_4) \Gamma(\frac{D}{2}-1-s_2) \Gamma(D-2-s_1-s_4)}{(4\pi)^{D/2} \Gamma(1+s_2) \Gamma(2-\frac{D}{2}+s_1+s_4) \Gamma(\frac{3D}{2}-3-s_1-s_2-s_4)} \\
&\quad \times \frac{1}{[(q_3 + q_4)^2]^{3-D+s_1+s_2+s_4}}. \tag{A6}
\end{aligned}$$

And then, we integrate out q_3 :

$$\begin{aligned}
I_{q_3} &\equiv \int \frac{d^D q_3}{(2\pi)^D} \frac{1}{(q_3^2)^{1+s_3} [(q_3 + q_4)^2]^{3-D+s_1+s_2+s_4}} \\
&= \frac{i(-)^{D/2} \Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 s_i) \Gamma(\frac{D}{2}-1-s_3) \Gamma(\frac{3D}{2}-3-s_1-s_2-s_4)}{(4\pi)^{D/2} \Gamma(1+s_3) \Gamma(3-D+s_1+s_2+s_4) \Gamma(2D-4-\sum_{i=1}^4 s_i)} \\
&\quad \times \frac{1}{(q_4^2)^{4-\frac{3D}{2}+\sum_{i=1}^4 s_i}}. \tag{A7}
\end{aligned}$$

Last, we integrate out q_4 :

$$\begin{aligned}
I_{q_4} &\equiv \int \frac{d^D q_4}{(2\pi)^D} \frac{1}{(q_4^2 - m_5^2) (q_4^2)^{4-\frac{3D}{2}+\sum_{i=1}^4 s_i}} \\
&= \frac{i}{(4\pi)^{D/2}} (-)^{5-\frac{3D}{2}+\sum_{i=1}^4 s_i} \left(\frac{1}{m_5^2}\right)^{5-2D+\sum_{i=1}^4 s_i} \Gamma(5-2D+\sum_{i=1}^4 s_i) \Gamma(2D-4-\sum_{i=1}^4 s_i). \tag{A8}
\end{aligned}$$

Together with Eqs. (A5-A8), one can have

$$\begin{aligned}
I_q &= \frac{-1}{(4\pi)^{2D}} (-)^{\sum_{i=1}^4 s_i} \left(\frac{1}{m_5^2}\right)^{5-2D+\sum_{i=1}^4 s_i} \left[\prod_{i=1}^4 \Gamma\left(\frac{D}{2}-1-s_i\right) \Gamma(1+s_i)^{-1} \right] \\
&\quad \times \Gamma\left(4-\frac{3D}{2}+\sum_{i=1}^4 s_i\right) \Gamma\left(5-2D+\sum_{i=1}^4 s_i\right). \tag{A9}
\end{aligned}$$

APPENDIX B: THE HYPERGEOMETRIC SERIES SOLUTIONS OF THE INTEGER LATTICE \mathbf{B}_{1345}

According the basis of integer lattice \mathbf{B}_{1345} , one can also construct GKZ hypergeometric series solutions below.

- $I_2 = [2, 5, \dots, 8, 10]$, i.e. the implement $J_2 = [1, \dots, 10] \setminus I_2 = [1, 3, 4, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(2)} = y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_4^{\frac{D}{2}-2} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(2)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(2)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B1})$$

- $I_3 = [2, 4, 6, 7, 9, 10]$, i.e. the implement $J_3 = [1, \dots, 10] \setminus I_3 = [1, 3, 5, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(3)} = y_1^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-2} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(3)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(3)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B2})$$

- $I_4 = [2, 4, \dots, 7, 10]$, i.e. the implement $J_4 = [1, \dots, 10] \setminus I_4 = [1, 3, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(4)} = y_1^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(4)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(4)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B3})$$

- $I_5 = [2, 3, 6, 8, 9, 10]$, i.e. the implement $J_5 = [1, \dots, 10] \setminus I_5 = [1, 4, 5, 7]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(5)} = y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-2} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(5)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(5)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B4})$$

- $I_6 = [2, 3, 5, 6, 8, 10]$, i.e. the implement $J_6 = [1, \dots, 10] \setminus I_6 = [1, 4, 7, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(6)} = y_2^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(6)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(6)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B5})$$

- $I_7 = [2, 3, 4, 6, 9, 10]$, i.e. the implement $J_7 = [1, \dots, 10] \setminus I_7 = [1, 5, 7, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(7)} = y_3^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(7)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(7)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B6})$$

- $I_8 = [2, 3, 4, 5, 6, 10]$, i.e. the implement $J_8 = [1, \dots, 10] \setminus I_8 = [1, 7, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(8)} = y_4^{\frac{3D}{2}-4} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(8)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(8)}(\mathbf{n}) = \frac{\Gamma(3 - D + \sum_{i=1}^4 n_i) \Gamma(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B7})$$

- $I_9 = [1, 6, \dots, 10]$, i.e. the implement $J_9 = [1, \dots, 10] \setminus I_9 = [2, 3, 4, 5]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(9)} = y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-2} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(9)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(9)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B8})$$

- $I_{10} = [1, 5, \dots, 8, 10]$, i.e. the implement $J_{10} = [1, \dots, 10] \setminus I_{10} = [2, 3, 4, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(10)} = y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(10)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(10)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B9})$$

- $I_{11} = [1, 4, 6, 7, 9, 10]$, i.e. the implement $J_{11} = [1, \dots, 10] \setminus I_{11} = [2, 3, 5, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(11)} = y_1^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(11)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(11)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B10})$$

- $I_{12} = [1, 4, \dots, 7, 10]$, i.e. the implement $J_{12} = [1, \dots, 10] \setminus I_{12} = [2, 3, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(12)} = y_1^{\frac{D}{2}-1} y_4^{\frac{3D}{2}-4} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(12)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(12)}(\mathbf{n}) = \frac{\Gamma(3 - D + \sum_{i=1}^4 n_i) \Gamma(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} \quad (\text{B11})$$

- $I_{13} = [1, 3, 6, 8, 9, 10]$, i.e. the implement $J_{13} = [1, \dots, 10] \setminus I_{13} = [2, 4, 5, 7]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(13)} = y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{D-3} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(13)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(13)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{B12})$$

- $I_{14} = [1, 3, 5, 6, 8, 10]$, i.e. the implement $J_{14} = [1, \dots, 10] \setminus I_{14} = [2, 4, 7, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(14)} = y_2^{\frac{D}{2}-1} y_4^{\frac{3D}{2}-4} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(14)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(14)}(\mathbf{n}) = \frac{\Gamma(3 - D + \sum_{i=1}^4 n_i) \Gamma(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{B13})$$

- $I_{15} = [1, 3, 4, 6, 9, 10]$, i.e. the implement $J_{15} = [1, \dots, 10] \setminus I_{15} = [2, 5, 7, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[1345]}^{(15)} = y_3^{\frac{D}{2}-1} y_4^{\frac{3D}{2}-4} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(15)}(\mathbf{n}) f_{[1345]} ,$$

$$c_{[1345]}^{(15)}(\mathbf{n}) = \frac{\Gamma(3-D+\sum_{i=1}^4 n_i)\Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i!\right]\Gamma(2-\frac{D}{2}+n_1)\Gamma(2-\frac{D}{2}+n_2)\Gamma(2-\frac{D}{2}+n_3)\Gamma(\frac{D}{2}+n_4)} . \quad (\text{B14})$$

- $I_{16} = [1, 3, 4, 5, 6, 10]$, i.e. the implement $J_{16} = [1, \dots, 10] \setminus I_{16} = [2, 7, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\begin{aligned} \Phi_{[1345]}^{(16)} &= y_4^{2D-5} \sum_{\mathbf{n}=0}^{\infty} c_{[1345]}^{(16)}(\mathbf{n}) f_{[1345]} , \\ c_{[1345]}^{(16)}(\mathbf{n}) &= \frac{\Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 n_i)\Gamma(5-2D+\sum_{i=1}^4 n_i)}{\prod_{i=1}^4 n_i! \Gamma(2-\frac{D}{2}+n_i)} . \end{aligned} \quad (\text{B15})$$

APPENDIX C: THE HYPERGEOMETRIC SERIES SOLUTIONS OF THE INTEGER LATTICE $\mathbf{B}_{\bar{1}345}$

According the basis of integer lattice $\mathbf{B}_{\bar{1}345}$, one can construct GKZ hypergeometric series solutions below.

- $I_1 = [1, 2, 7, \dots, 10]$, i.e. the implement $J_1 = [1, \dots, 10] \setminus I_1 = [3, 4, 5, 6]$. The corresponding hypergeometric series solution is written as

$$\begin{aligned} \Phi_{[\bar{1}345]}^{(1)} &= y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(1)}(\mathbf{n}) f_{[\bar{1}345]} , \\ f_{[\bar{1}345]} &= (y_4)^{n_1} (y_1)^{n_2} (y_2)^{n_3} (y_3)^{n_4} , \end{aligned} \quad (\text{C1})$$

where the coefficient is

$$c_{[\bar{1}345]}^{(1)}(\mathbf{n}) = \frac{\Gamma(\frac{D}{2}+\sum_{i=1}^4 n_i)\Gamma(1+\sum_{i=1}^4 n_i)}{\prod_{i=1}^4 n_i! \Gamma(\frac{D}{2}+n_i)} . \quad (\text{C2})$$

- $I_2 = [1, 2, 6, \dots, 9]$, i.e. the implement $J_2 = [1, \dots, 10] \setminus I_2 = [3, 4, 5, 10]$. The corresponding hypergeometric series solution is written as

$$\begin{aligned} \Phi_{[\bar{1}345]}^{(2)} &= y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(2)}(\mathbf{n}) f_{[\bar{1}345]} , \\ c_{[\bar{1}345]}^{(2)}(\mathbf{n}) &= \frac{\Gamma(1+\sum_{i=1}^4 n_i)\Gamma(2-\frac{D}{2}+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i!\right]\Gamma(2-\frac{D}{2}+n_1)\Gamma(\frac{D}{2}+n_2)\Gamma(\frac{D}{2}+n_3)\Gamma(\frac{D}{2}+n_4)} . \end{aligned} \quad (\text{C3})$$

- $I_3 = [1, 2, 5, 7, 8, 10]$, i.e. the implement $J_3 = [1, \dots, 10] \setminus I_3 = [3, 4, 6, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(3)} = y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(3)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(3)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (C4)$$

- $I_4 = [1, 2, 5, \dots, 8]$, i.e. the implement $J_4 = [1, \dots, 10] \setminus I_4 = [3, 4, 9, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(4)} = y_1^{\frac{D}{2}-1} y_2^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(4)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(4)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (C5)$$

- $I_5 = [1, 2, 4, 7, 9, 10]$, i.e. the implement $J_5 = [1, \dots, 10] \setminus I_5 = [3, 5, 6, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(5)} = y_1^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(5)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(5)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (C6)$$

- $I_6 = [1, 2, 4, 6, 7, 9]$, i.e. the implement $J_6 = [1, \dots, 10] \setminus I_6 = [3, 5, 8, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(6)} = y_1^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(6)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(6)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (C7)$$

- $I_7 = [1, 2, 4, 5, 7, 10]$, i.e. the implement $J_7 = [1, \dots, 10] \setminus I_7 = [3, 6, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(7)} = y_1^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(7)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(7)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{C8})$$

- $I_8 = [1, 2, 4, 5, 6, 7]$, i.e. the implement $J_8 = [1, \dots, 10] \setminus I_8 = [3, 8, 9, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(8)} = y_1^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(8)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(8)}(\mathbf{n}) = \frac{\Gamma(3 - D + \sum_{i=1}^4 n_i) \Gamma(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} + n_2) \Gamma(2 - \frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{C9})$$

- $I_9 = [1, 2, 3, 8, 9, 10]$, i.e. the implement $J_9 = [1, \dots, 10] \setminus I_9 = [4, 5, 6, 7]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(9)} = y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(9)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(9)}(\mathbf{n}) = \frac{\Gamma(1 + \sum_{i=1}^4 n_i) \Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{C10})$$

- $I_{10} = [1, 2, 3, 6, 8, 9]$, i.e. the implement $J_{10} = [1, \dots, 10] \setminus I_{10} = [4, 5, 7, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(10)} = y_2^{\frac{D}{2}-1} y_3^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(10)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(10)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(\frac{D}{2} + n_4)} . \quad (\text{C11})$$

- $I_{11} = [1, 2, 3, 5, 8, 10]$, i.e. the implement $J_{11} = [1, \dots, 10] \setminus I_{11} = [4, 6, 7, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(11)} = y_2^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(11)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(11)}(\mathbf{n}) = \frac{\Gamma(2 - \frac{D}{2} + \sum_{i=1}^4 n_i) \Gamma(3 - D + \sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} + n_3) \Gamma(2 - \frac{D}{2} + n_4)} . \quad (\text{C12})$$

- $I_{12} = [1, 2, 3, 5, 6, 8]$, i.e. the implement $J_{12} = [1, \dots, 10] \setminus I_{12} = [4, 7, 9, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(12)} = y_2^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(12)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(12)}(\mathbf{n}) = \frac{\Gamma(3-D+\sum_{i=1}^4 n_i) \Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2-\frac{D}{2}+n_1) \Gamma(2-\frac{D}{2}+n_2) \Gamma(\frac{D}{2}+n_3) \Gamma(2-\frac{D}{2}+n_4)} . \quad (\text{C13})$$

- $I_{13} = [1, 2, 3, 4, 9, 10]$, i.e. the implement $J_{13} = [1, \dots, 10] \setminus I_{13} = [5, 6, 7, 8]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(13)} = y_3^{\frac{D}{2}-1} y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(13)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(13)}(\mathbf{n}) = \frac{\Gamma(2-\frac{D}{2}+\sum_{i=1}^4 n_i) \Gamma(3-D+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2}+n_1) \Gamma(2-\frac{D}{2}+n_2) \Gamma(2-\frac{D}{2}+n_3) \Gamma(\frac{D}{2}+n_4)} . \quad (\text{C14})$$

- $I_{14} = [1, 2, 3, 4, 6, 9]$, i.e. the implement $J_{14} = [1, \dots, 10] \setminus I_{14} = [5, 7, 8, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(14)} = y_3^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(14)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(14)}(\mathbf{n}) = \frac{\Gamma(3-D+\sum_{i=1}^4 n_i) \Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(2-\frac{D}{2}+n_1) \Gamma(2-\frac{D}{2}+n_2) \Gamma(2-\frac{D}{2}+n_3) \Gamma(\frac{D}{2}+n_4)} . \quad (\text{C15})$$

- $I_{15} = [1, 2, 3, 4, 5, 10]$, i.e. the implement $J_{15} = [1, \dots, 10] \setminus I_{15} = [6, 7, 8, 9]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(15)} = y_4^{\frac{D}{2}-1} \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(15)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(15)}(\mathbf{n}) = \frac{\Gamma(3-D+\sum_{i=1}^4 n_i) \Gamma(4-\frac{3D}{2}+\sum_{i=1}^4 n_i)}{\left[\prod_{i=1}^4 n_i! \right] \Gamma(\frac{D}{2}+n_1) \Gamma(2-\frac{D}{2}+n_2) \Gamma(2-\frac{D}{2}+n_3) \Gamma(2-\frac{D}{2}+n_4)} . \quad (\text{C16})$$

- $I_{16} = [1, 2, 3, 4, 5, 6]$, i.e. the implement $J_{16} = [1, \dots, 10] \setminus I_{16} = [7, 8, 9, 10]$. The corresponding hypergeometric series solution is written as

$$\Phi_{[\bar{1}345]}^{(16)} = \sum_{\mathbf{n}=0}^{\infty} c_{[\bar{1}345]}^{(16)}(\mathbf{n}) f_{[\bar{1}345]} ,$$

$$c_{[\bar{1}345]}^{(16)}(\mathbf{n}) = \frac{\Gamma(4 - \frac{3D}{2} + \sum_{i=1}^4 n_i) \Gamma(5 - 2D + \sum_{i=1}^4 n_i)}{\prod_{i=1}^4 n_i! \Gamma(2 - \frac{D}{2} + n_i)} . \quad (\text{C17})$$

- [1] L. Linssen et al., Physics and Detectors at CLIC: CLIC Conceptual Design Report, CERN-2012-003, arXiv:1202.5940 [physics.ins-det].
- [2] T. Behnke et al., The International Linear Collider Technical Design Report - Volume I: Executive Summary, arXiv:1306.6327 [physics.acc-ph].
- [3] J.B.G. da Costa et al. (CEPC Study Group), CEPC Conceptual Design Report: Volume 2 - Physics & Detector, IHEP-CEPC-DR-2018-02, arXiv:1811.10545 [hep-ex].
- [4] A. Abada et al. (FCC Collaboration), FCC Physics Opportunities: Future Circular Collider Conceptual Design Report Volume 1, Eur. Phys. J. C 79 (2019) 474.
- [5] I.B. Alonso et al., High-Luminosity Large Hadron Collider (HL-LHC): Technical design report, CERN-2020-010.
- [6] G. Heinrich, Collider Physics at the Precision Frontier, Phys. Rept. 922 (2021) 1-69.
- [7] V.A. Smirnov, Applied asymptotic expansions in momenta and masses, Springer, Berlin 2002.
- [8] M. Misiak, M. Münz, Two loop mixing of dimension five flavor changing operators, Phys. Lett. B 344 (1995) 308.
- [9] Y. Schröder, Tackling the infrared problem of thermal QCD, Nucl. Phys. (Proc. Suppl.) 129 (2004) 572.
- [10] G. t'Hooft, M. Veltman, Nucl. Phys. B 153 (1979) 365.
- [11] G. Passarino, M. Veltman, Nucl. Phys. B 160 (1979) 151.
- [12] A. Denner, Techniques for calculation of electroweak radiative corrections at the one loop level and results for W physics at LEP-200, Fortschr. Phys. 41 (1993) 307-420.
- [13] V.A. Smirnov, Analytic Tools for Feynman Integrals, Springer, Heidelberg, 2012, Springer Tracts Mod. Phys. 250 (2012) 1-296.
- [14] C. Ford, I. Jack, D.R.T. Jones, The Standard model effective potential at two loops, Nucl. Phys. B 387 (1992) 373-390, [Erratum-ibid. 504 (1997) 551-552].
- [15] A.I. Davydychev, J.B. Tausk, Two loop selfenergy diagrams with different masses and the momentum expansion, Nucl. Phys. B 397 (1993) 123.

- [16] A.I. Davydychev, V.A. Smirnov, J.B. Tausk, Large momentum expansion of two-loop self-energy diagrams with arbitrary masses, Nucl. Phys. B 410 (1993) 325.
- [17] R. Scharf, J.B. Tausk, Scalar two loop integrals for gauge boson selfenergy diagrams with a massless fermion loop, Nucl. Phys. B 412 (1994) 523.
- [18] J.R. Espinosa, R.J. Zhang, Complete two-loop dominant corrections to the mass of the lightest CP-even Higgs boson in the minimal supersymmetric standard model, Nucl. Phys. B 586 (2000) 3.
- [19] D.J. Broadhurst, Three loop on-shell charge renormalization without integration: Lambda-MS (QED) to four loops, Z. Phys. C 54 (1992) 599.
- [20] S. Laporta, E. Remiddi, The Analytical value of the electron light-light graphs contribution to the muon (g-2) in QED, Phys. Lett. B 301 (1993) 440.
- [21] L. Avdeev, J. Fleischer, S. Mikhailov, O. Tarasov, $\mathcal{O}(\alpha\alpha_s^2)$ correction to the electroweak ρ parameter, Phys. Lett. B 336 (1994) 560, [Erratum-ibid. 349 (1995) 597].
- [22] J. Fleischer, O.V. Tarasov, Application of conformal mapping and Padé approximants ($\omega P's$) to the calculation of various two-loop Feynman diagrams, Nucl. Phys. B, Proc. Suppl. 37 (1994) 115.
- [23] L.V. Avdeev, Recurrence relations for three-loop prototypes of bubble diagrams with a mass, Comput. Phys. Commun. 98 (1996) 15.
- [24] J. Fleischer, M.Y. Kalmykov, Single mass scale diagrams: Construction of a basis for the ϵ -expansion, Phys. Lett. B 470 (1999) 168.
- [25] D.J. Broadhurst, Massive three-loop Feynman diagrams reducible to SC* primitives of algebras of the sixth root of unity, Eur. Phys. J. C 8 (1999) 311.
- [26] K.G. Chetyrkin, M. Steinhauser, The Relation between the MS-bar and the on-shell quark mass at order α_s^2 , Nucl. Phys. B 573 (2000) 617.
- [27] A.I. Davydychev, M.Y. Kalmykov, Massive Feynman diagrams and inverse binomial sums, Nucl. Phys. B 699 (2004) 3.
- [28] Y. Schröder, A. Vuorinen, High-precision ϵ -expansions of single-mass-scale four-loop vacuum bubbles, JHEP 06 (2005) 051.
- [29] M.Y. Kalmykov, About higher order ϵ -expansion of some massive two- and three-loop master-integrals, Nucl. Phys. B 718 (2005) 276.

- [30] M.Y. Kalmykov, Gauss hypergeometric function: Reduction, ϵ -expansion for integer/half-integer parameters and Feynman diagrams, JHEP 04 (2006) 056.
- [31] S. Bekavac, A.G. Grozin, D. Seidel, V.A. Smirnov, Three-loop on-shell Feynman integrals with two masses, Nucl. Phys. B 819 (2009) 183.
- [32] V.V. Bytev, M.Y. Kalmykov, B.A. Kniehl, Differential reduction of generalized hypergeometric functions from Feynman diagrams: One-variable case, Nucl. Phys. B 836 (2010) 129.
- [33] J. Grigo, J. Hoff, P. Marquard, M. Steinhauser, Moments of heavy quark correlators with two masses: exact mass dependence to three loops, Nucl. Phys. B 864 (2012) 580.
- [34] V.V. Bytev, M.Y. Kalmykov, B.A. Kniehl, HYPERDIRE, HYPERgeometric functions Differential REduction: MATHEMATICA-based packages for differential reduction of generalized hypergeometric functions ${}_pF_{p-1}$, F_1 , F_2 , F_3 , F_4 , Comput. Phys. Commun. 184 (2013) 2332.
- [35] A. Freitas, Three-loop vacuum integrals with arbitrary masses, JHEP 11 (2016) 145.
- [36] S.P. Martin, D.G. Robertson, Evaluation of the general 3-loop vacuum Feynman integral, Phys. Rev. D 95 (2017) 016008.
- [37] S.P. Martin, Effective potential at three loops, Phys. Rev. D 96 (2017) 096005.
- [38] Z.-H. Gu, H.-B. Zhang, Three-loop vacuum integral with four-propagators using hypergeometry, Chin. Phys. C 43 (2019) 083102.
- [39] Z.-H. Gu, H.-B. Zhang, T.-F. Feng, Hypergeometric expression for a three-loop vacuum integral, Int. J. Mod. Phys. A 35 (2020) 2050089.
- [40] H.-B. Zhang, T.-F. Feng, GKZ hypergeometric systems of the three-loop vacuum Feynman integrals, JHEP 05 (2023) 075.
- [41] Y. Schröder, A. Vuorinen, High-precision epsilon expansions of single-mass-scale four-loop vacuum bubbles, JHEP 06 (2005) 051.
- [42] E. Bejdakic, Y. Schröder, Hypergeometric representation of a four-loop vacuum bubble, Nucl. Phys. B (Proc. Suppl.) 160 (2006) 155-159.
- [43] S. Laporta, High-precision epsilon expansions of massive four-loop vacuum bubbles, Phys. Lett. B 549 (2002) 115.
- [44] Y. Schröder, Automatic reduction of four-loop bubbles, Nucl. Phys. B (Proc. Suppl.) 116 (2003) 402-406.

- [45] K. Kajantie, M. Laine, Y. Schröder, Simple way to generate high order vacuum graphs, *Phys. Rev. D* 65 (2002) 045008.
- [46] X. Liu, Y.-Q. Ma, Determining arbitrary Feynman integrals by vacuum integrals, *Phys. Rev. D* 99 (2019) 071501(R).
- [47] X. Liu, Y.-Q. Ma, AMFlow: A Mathematica package for Feynman integrals computation via auxiliary mass flow, *Comput. Phys. Commun.* 283 (2023) 108565.
- [48] T. Regge, Algebraic Topology Methods in the Theory of Feynman Relativistic Amplitudes, In: *Battelle Rencontres - 1967 Lectures in Math. and Phys.*, edited by C. M. DeWitt and J. A. Wheeler (W. A. Benjamin, Inc., New York, 1967), pp. 433-458.
- [49] E.E. Boos, A.I. Davydychev, A method for calculating vertex-type Feynman integrals, *Vestn.Mosk.Univ.Fiz.Astron.* 28N3 (1987) 8-12.
- [50] E.E. Boos, A.I. Davydychev, A Method of evaluating massive Feynman integrals, *Theor. Math. Phys.* 89 (1991) 1052.
- [51] A.I. Davydychev, Some exact results for N point massive Feynman integrals, *J. Math. Phys.* 32 (1991) 1052.
- [52] A.I. Davydychev, Recursive algorithm of evaluating vertex type Feynman integrals, *J. Phys. A* 25 (1992) 5587.
- [53] A.I. Davydychev, General results for massive N point Feynman diagrams with different masses, *J. Math. Phys.* 33 (1992) 358.
- [54] N.I. Ussyukina, A.I. Davydychev, An Approach to the evaluation of three and four point ladder diagrams, *Phys. Lett. B* 298 (1993) 363.
- [55] F.A. Berends, M. Böhm, M. Buza, R. Scharf, Closed expressions for specific massive multiloop self-energy integrals, *Z. Phys. C* 63 (1994) 227.
- [56] V.A. Smirnov, Analytical result for dimensionally regularized massless on shell double box, *Phys. Lett. B* 460 (1999) 397-404.
- [57] J.B. Tausk, Nonplanar massless two loop Feynman diagrams with four on-shell legs, *Phys. Lett. B* 469 (1999) 225-234.
- [58] A.I. Davydychev, Explicit results for all orders of the epsilon expansion of certain massive and massless diagrams, *Phys. Rev. D* 61 (2000) 087701.
- [59] O.V. Tarasov, Application and explicit solution of recurrence relations with respect to space-

- time dimension, Nucl. Phys. B, Proc. Suppl. 89 (2000) 237.
- [60] J. Fleischer, F. Jegerlehner, O. V. Tarasov, A New hypergeometric representation of one loop scalar integrals in d dimensions, Nucl. Phys. B 672 (2003) 303.
 - [61] A.I. Davydychev, Geometrical methods in loop calculations and the three-point function, Nucl. Instrum. Meth. A 559 (2006) 293.
 - [62] M.Y. Kalmykov, B.A. Kniehl, Towards all-order Laurent expansion of generalized hypergeometric functions around rational values of parameters, Nucl. Phys. B 809 (2009) 365.
 - [63] M.Y. Kalmykov, B.A. Kniehl, Counting master integrals: Integration by parts versus differential reduction, Phys. Lett. B 702 (2011) 268.
 - [64] M.Y. Kalmykov, B.A. Kniehl, Mellin-Barnes representations of Feynman diagrams, linear systems of differential equations, and polynomial solutions, Phys. Lett. B 714 (2012) 103.
 - [65] V. Bytev, M. Kalmykov, HYPERDIRE hypergeometric functions differential reduction: Mathematica-based packages for the differential reduction of generalized hypergeometric function: Horn-type hypergeometric functions of two variables, Comput. Phys. Commun. 189 (2015) 128.
 - [66] V.V. Bytev, M.Y. Kalmykov, HYPERDIRE, HYPERgeometric functions Differential Reduction: Mathematica-based packages for the differential reduction of generalized hypergeometric functions: Lauricella function F_c of three variables, Comput. Phys. Commun. 206 (2016) 78.
 - [67] M.Y. Kalmykov, B.A. Kniehl, Counting the number of master integrals for sunrise diagrams via the Mellin-Barnes representation, JHEP 07 (2017) 031.
 - [68] T.-F. Feng, C.-H. Chang, J.-B. Chen, Z.-H. Gu, H.-B. Zhang, Evaluating Feynman integrals by the hypergeometry, Nucl. Phys. B 927 (2018) 516.
 - [69] T.-F. Feng, C.-H. Chang, J.-B. Chen, H.-B. Zhang, The system of partial differential equations for the C_0 function, Nucl. Phys. B 940 (2019) 130.
 - [70] S. Abreu, R. Britto, C. Duhr, E. Gardi, J. Matthew, From positive geometries to a coaction on hypergeometric functions, JHEP 02 (2020) 122.
 - [71] B. Ananthanarayan, S. Banik, S. Ghosh, New series representations for the two-loop massive sunset diagram, Eur. Phys. J. C 80 (2020) 606.
 - [72] B. Ananthanarayan, S. Banik, S. Friot, S. Ghosh, Multiple Series Representations of N -fold

- Mellin-Barnes Integrals, Phys. Rev. Lett. 127 (2021) 151601, arXiv:2012.15108 [hep-th].
- [73] E. Nasrollahpoursamami, Periods of Feynman Diagrams and GKZ D-modules, arXiv:1605.04970 [math-ph].
 - [74] I.M. Gel'fand, General theory of hypergeometric functions, Soviet Math. Dokl. 33 (1986) 573.
 - [75] I.M. Gel'fand, M.I. Graev, A.V. Zelevinsky, Holonomic systems of equations and series of hypergeometric type, Soviet Math. Dokl. 36 (1988) 5.
 - [76] I.M. Gel'fand, A.V. Zelevinsky, M. M. Kapranov, Hypergeometric functions and toral manifold, Soviet Math. Dokl. 37 (1988) 678.
 - [77] I.M. Gel'fand, M.M. Kapranov, and A.V. Zelevinsky, Generalized Euler integrals and A-hypergeometric systems, Adv. in Math. 84 (1990) 255.
 - [78] I. M. Gelfand, A. V. Zelevinskii, and M. M. kapranov, Hypergeometric functions and toric varieties, Functional. Anal. Appl. 23 (1989) 94-106.
 - [79] L. Cruz, Feynman integrals as A-hypergeometric functions, JHEP 12 (2019) 123.
 - [80] R. Klausen, Hypergeometric Series Representations of Feynman integrals by GKZ Hypergeometric Systems, JHEP 04 (2020) 121.
 - [81] R.N. Lee, A.A. Pomeransky, Critical points and number of master integrals, JHEP 11 (2013) 165.
 - [82] T. Oaku, Algorithms for Functions, restrictions, and algebraic local cohomology groups of D-Modules, Adv. Appl. Math. 19 (1997) 61.
 - [83] U. Walther, Algorithmic computation of local cohomology modules and the local cohomological dimension of algebraic varieties, J. Pure Appl. Algebra 139 (1999) 303.
 - [84] T. Oaku, N. Takayama, Algorithms for D-modules - restriction, tensor product, localization, and local cohomology groups, J. Pure Appl. Algebra 156 (2001) 267.
 - [85] T.-F. Feng, C.-H. Chang, J.-B. Chen, H.-B. Zhang, GKZ-hypergeometric systems for Feynman integrals, Nucl. Phys. B 953 (2020) 114952, arXiv:1912.01726.
 - [86] T.-F. Feng, H.-B. Zhang, Y.-Q. Dong, Y. Zhou, GKZ-system of the 2-loop self energy with 4 propagators, Eur. Phys. J. C 83 (2023) 314, arXiv:2209.15194.
 - [87] T.-F. Feng, H.-B. Zhang, C.-H. Chang. Feynman integrals of Grassmannians, Phys. Rev. D 106 (2022) 116025, arXiv:2206.04224.
 - [88] W. Miller Jr., Lie theory and the hypergeometric functions, J. Math. Mech. 17 (1968) 1143.

- [89] W. Miller Jr., Lie theory and generalized hypergeometric functions, *SIAM. J. Math. Anal.* 3 (1972) 31.
- [90] F. Loebbert, D. Müller, H. Münkler, Yangian Bootstrap for Conformal Feynman Integrals, *Phys. Rev. D* 101 (2020) 066006.
- [91] A. Klemm, C. Nega, and R. Safari, The l-loop Banana Amplitude from GKZ Systems and relative Calabi-Yau Periods, *JHEP* 04 (2020) 088.
- [92] K. Bönisch, F. Fischbach, A. Klemm, C. Nega, R. Safari, Analytic Structure of all Loop Banana Amplitudes, *JHEP* 05 (2021) 066.
- [93] M. Hidding, DiffExp, a Mathematica package for computing Feynman integrals in terms of one-dimensional series expansions, *Comput. Phys. Commun.* 269 (2021) 108125.
- [94] M. Borinsky, Tropical Monte Carlo quadrature for Feynman integrals, *Ann. Inst. H. Poincaré Comb. Phys. Interact.* 10 (2023) 635-685, arXiv: 2008.12310 [math-ph].
- [95] M. Kalmykov, V. Bytev, B. Kniehl, S.-O. Moch, B. Ward, S. Yost, Hypergeometric Functions and Feynman Diagrams, arXiv:2012.14492 [hep-th].
- [96] F. Tellander, M. Helmer, Cohen-Macaulay Property of Feynman Integrals, *Commun. Math. Phys.* 399 (2023) 1021-1037.
- [97] R.P. Klausen, Kinematic singularities of Feynman integrals and principal A-determinants, *JHEP* 02 (2022) 004.
- [98] S. Mizera, and S. Telen, Landau Discriminants, *JHEP* 08 (2022) 200.
- [99] N. Arkani-Hamed, A. Hillman, S. Mizera, Feynman Polytopes and the tropical geometry of UV and IR divergence, *Phys. Rev. D* 105 (2022) 125013.
- [100] V. Chestnov, F. Gasparotto, M. K. Mandal, P. Mastrolia, S. J. Matsubara-Heo, H. J. Munch, N. Takayamac, Macaulay Matrix for Feynman Integrals: Linear Relations and Intersection Numbers, *JHEP* 09 (2022) 187.
- [101] U. Walther, On Feynman graphs, matroids, and GKZ-systems, *Lett. Math. Phys.* 112 (2022) 120.
- [102] H. J. Munch, Feynman Integral Relations from GKZ Hypergeometric Systems, *PoS LL2022* (2022) 042.
- [103] B. Ananthanarayan, S. Banik, S. Bera, S. Datta, FeynGKZ: a Mathematica package for solving Feynman integrals using GKZ hypergeometric systems, *Comput. Phys. Commun.*

287 (2023) 108699.

- [104] I. Dubovyk, J. Gluza, G. Somogyi, Mellin-Barnes Integrals: A Primer on Particle Physics Applications, arXiv:2211.13733 [hep-ph].
- [105] V. Chestnov, S. J. Matsubara-Heo, H. J. Munch, N. Takayama, Restrictions of Pfaffian systems for Feynman integrals, JHEP 11 (2023) 202.
- [106] R. P. Klausen, Hypergeometric feynman integrals, arXiv:2302.13184 [hep-th].
- [107] F. Caloro, P. McFadden, A-hypergeometric functions and creation operators for Feynman and Witten diagrams, arXiv:2309.15895 [hep-th].
- [108] H. J. Munch, Evaluating Feynman Integrals Using D-modules and Tropical Geometry, arXiv:2401.00891 [hep-th].
- [109] A.V. Smirnov, N.D. Shapurov, L.I. Vysotsky, FIESTA5: Numerical highperformance Feynman integral evaluation, Comput. Phys. Commun. 277 (2022) 108386.